arXiv:1205.3940v2 [math.LO] 16 Jul 2012

New Directions in Categorical Logic, for Classical, Probabilistic and Quantum Logic

Bart Jacobs

Institute for Computing and Information Sciences (iCIS),
Radboud University Nijmegen, The Netherlands.
Web address: `www.cs.ru.nl/B.Jacobs`

March 4, 2013

Abstract. Traditionally in categorical logic predicates on an object/type X are represented as subobjects of X. Here we break with that tradition and use maps of the form $p\colon X \to X + X$ with $[\mathrm{id}, \mathrm{id}] \circ p = \mathrm{id}$ as predicates. This new view gives a more dynamic, measurement-oriented view on predicates, that works well especially in a quantitative setting. In classical logic (in the category of sets) these new predicates coincide with the traditional ones (subsets, or characteristic maps $X \to \{0,1\}$); in probabilistic logic (in the category of sets and stochastic matrices), the new predicates correspond to fuzzy predicates $X \to [0,1]$; and in quantum logic (in Hilbert spaces) they correspond to effects (positive endomaps below the identity), which may be understood as fuzzy predicates on a changed basis.

It is shown that, under certain conditions about coproducts $+$, predicates $p\colon X \to X + X$ form effect algebras and carry a scalar multiplication (with probabilities). Suitable substitution functors give rise to indexed/fibred categories. In the quantum case the famous Born rule — describing the probability of observation outcomes — follows directly from the form of these substitution functors: probability calculation becomes substitution in predicate logic. Moreover, the characteristic maps associated with predicates provide tests in a dynamic logic, and turn out to capture measurement in a form that uniformly covers the classical, probabilistic and quantum case. The probabilities incorporated in predicates (as eigenvalues) serves as weights for the possible measurement outcomes.

1 Introduction

Mathematical logic started in the 19th century with George Boole's *Laws of Thought*. Since then 'Boolean algebras' form the (algebraic) basis of formal logic. In the 20th century these Boolean algebras have been generalised, mainly in two directions, see Figure 1. Of these two directions the one on the left is most familiar and influential. It has given rise to the edifice of topos theory, where the logic naturally associated with geometric structures is intuitionistic — in the sense that the double-negation law $\neg\neg P = P$ fails. Also, it has given rise to Martin-Löf's intuitionistic type theory [47,52], which forms the foundation for the theorem prover Coq (and other such computerised formal systems). In this tradition it is claimed that the natural logic of computation is intuitionistic.



Fig. 1. Generalisations of Boolean logic

However, there are generalisations of Boolean logic where double-negation holds, like linear logic or quantum logic. In this tradition negation is often written as superscript $P^{\perp}$ and called orthocomplement. The logic of the quantum world has first been formalised by von Neumann and Birkhoff in terms of orthomodular lattices [50,6] (see also the monograph [40]). But these orthomodular lattices are rather awkward mathematical structures that have not led to a formal calculus of propositions (or predicates) capturing relevant logical phenomena in physics.

A second generalisation, started in the 1990s, involves so-called effect algebras [21] and effect modules, introduced in various forms (*e.g.* as D-posets [13] or as weak orthoalgebras in [23]) and by various people (see [20]), notably by theoretical physicists Foulis, Bennet, Gudder and Pulmannová, who published their work in the physics literature, largely outside the range of perception of traditional logicians. These effect algebras are partial commutative monoids with an orthocomplement $(-)^{\perp}$. Examples come from classical logic (Boolean algebras), quantum logic (orthomodular lattices, like the projections of a Hilbert space), probability theory (the unit interval $[0,1]$) and measure theory (step functions, taking finitely many values). Effect algebras form a common generalisation of structures from logic and probability theory, making it possible to see both negation $P \mapsto \neg P$ in logic and opposite probability $r \mapsto 1 - r$, as instances of the same orthocomplement operation. Partial commutative monoids have appeared in programming language semantics early on, notably in the partially additive structures of [3], and also in models of linear logic [26]. But orthocomplement is typical of the more restrictive quantum context.

Effect modules are effect algebras with an additional scalar multiplication, where the scalars are probabilities (usually from $[0,1]$). Intuitively, these effect modules can be seen as vector spaces, defined not over the real or complex numbers, but over $[0,1]$. They capture some essential properties of structures used in probability theory. In a quantum context the effects $\mathcal{E}f(H)$ of a Hilbert space H (the positive operators below the identity) form the main example, but more generally, $[0,1]$-valued functions form examples.

The basic theory of effect algebras and modules has been developed by several physicists who (understandably) lacked a background in modern logic, (program) semantics, and categorical logic. Especially the unfamiliarity with the modern methods of categorical logic seriously hampered, or even blocked, progress in this field. As a result these founders failed to identify certain fundamental relationships and similarities with programming semantics and logic (notably the categorical dualities between convex sets and effect modules [36], described in quantum theoretic terms as the duality between states and effects, or between Schrödinger's and Heisenberg's view, see [29, Chapter 2]).

The main topic of this paper however is different. It involves a natural representation of logical formulas in categories, supporting the right branch in Figure 1. The traditional approach in categorical logic of representing predicates as subobjects does not work, since double-negation only rarely holds for such predicates.

The essence of the novel approach that we propose here is to represent a predicate on a set/type/object X as a map:

$$X \xrightarrow{p} X+X \qquad \text{satisfying} \qquad \begin{tikzcd} X \arrow[r, "p"] \arrow[dr, equal] & X+X \arrow[d, "\nabla={[\mathrm{id},\mathrm{id}]}"] \\ & X \end{tikzcd} \tag{1}$$

where $+$ describes a coproduct in some category. This approach is certainly not as general as the subobject-based approach, but it does have some distinctive advantages. Most importantly, orthocomplement is easy, namely just post-compose a predicate $p\colon X \to X+X$ with the swap map $[\kappa_2,\kappa_1]\colon X+X \xrightarrow{\cong} X+X$. Thus $p^\perp = [\kappa_2,\kappa_1] \circ p$, immediately giving us $p^{\perp\perp} = p$. Notice that we can take the first coprojection $\kappa_1\colon X \to X+X$ as "true" and the second coprojection $\kappa_2\colon X \to X+X$ as "false". Orthocomplement $(-)^\perp$ exchanges true and false.

The representation (1) describes a predicate as a coalgebra, of the functor $Y \mapsto Y+Y$. Indeed, this representation emphasises the dynamic character of predicates, describing a transition to successor states. This is most clearly the case in measurement specified by a predicate, see Section 10.

We briefly describe this representation (1) for our main application areas: classical logic, probabilistic logic, and quantum logic. Details will appear later on in the paper.

- The world of *classical logic* appears when we consider a map $p\colon X \to X+X$ with $\nabla \circ p = \mathrm{id}$ in the category **Sets** of sets and functions. The coproduct $+$ in **Sets** is simply disjoint union. The equation $\nabla \circ p = \mathrm{id}$ means that $p(x) \in X+X$ can be only of the form $\kappa_1 x$, in the left component of $X+X$, or of the form $\kappa_2 x$, in the right component. Thus we can identify p with a subset $U \subseteq X$, namely $U = \{x \mid p(x) = \kappa_1 x\}$. Then $\neg U = \{x \mid p(x) = \kappa_2 x\} = \{x \mid p^\perp(x) = \kappa_1 x\}$. We shall see that this approach makes sense more generally, in so-called extensive categories (which include toposes).
- As (simple) *probabilistic* model we take a category of sets and stochastic relations between them. More formally, this is the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the discrete probability distribution monad $\mathcal{D}$ on **Sets**. This monad maps a set X to the collec-

tion of formal convex combinations of elements in X.

$$\mathcal{D}(X) = \{r_1 x_1 + \cdots + r_n x_n \mid x_i \in X, r_i \in [0,1] \text{ with } \sum_i r_i = 1\}.$$

An object of the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ is just a set, and a map $f\colon X \to Y$ is a function $f\colon X \to \mathcal{D}(Y)$; it can be seen as a stochastic relation between X and Y, or also as a Markov chain — with probability of a transition $x \to x_i$ described by $r_i \in [0,1]$ in the convex sum $f(x) = r_1 x_1 + \cdots + r_n x_n$. The identity $X \to X$ sends $x \in X$ to the singleton (or 'Dirac') distribution $1x \in \mathcal{D}(X)$. This category $\mathcal{K}\ell(\mathcal{D})$ has coproducts $+$ as in **Sets**. A predicate (1) is given by a function $p\colon X \to \mathcal{D}(X + X)$ and the equation $\nabla \circ p = \mathrm{id}$ in the Kleisli category means that each $p(x) \in \mathcal{D}(X + X)$ can only be a convex combination of the left and right elements $\kappa_1 x, \kappa_2 x \in X + X$. Thus we can write:

$$p(x) = \varphi(x)\kappa_1 x + \psi(x)\kappa_2 x \qquad \text{where} \qquad \begin{cases} \varphi(x), \psi(x) \in [0,1] \\ \text{satisfy } \varphi(x) + \psi(x) = 1. \end{cases}$$

Hence we may also write p as convex combination:

$$p(x) = \varphi(x)\kappa_1 x + (1 - \varphi(x))\kappa_2 x.$$

Thus, a predicate (1) on X in $\mathcal{K}\ell(\mathcal{D})$ can be identified with a "fuzzy" predicate $\varphi\colon X \to [0,1]$. In this way we see that fuzzy logic — where truth comes in degrees — naturally emerges from probabilistic modelling. The orthocomplement $\varphi^\perp$ corresponds to the function $x \mapsto 1 - \varphi(x)$. It involves pointwise the orthocomplement on the unit interval $[0,1]$. As we shall see, the structure that is naturally associated with fuzzy predicates is described in terms of effect algebras and modules.

- For the *quantum logic* perspective we turn to the category **Hilb** of Hilbert spaces and linear continuous (bounded) maps. The coproduct in **Hilb** is special, because it is at the same time a product. This is called a 'biproduct', written as $\oplus$. Thus, a predicate (1) on a Hilbert space X is given by a pair of maps $p = \langle p_1, p_2\rangle\colon X \to X \oplus X$. The condition $\nabla \circ p = \mathrm{id}$ means that $p_1 + p_2 = \mathrm{id}$, where $+$ now describes (pointwise) addition of maps. Thus, the predicate p is determined by its first part p_1, since $p_2 = \mathrm{id} - p_1$.
  The category **Hilb** is a so-called dagger category: there is a functor $(-)^\dagger\colon \mathbf{Hilb}^{\mathrm{op}} \to \mathbf{Hilb}$ that is the identity on objects and complex conjugation on morphisms. Thus $f^{\dagger\dagger} = f$ for a map f in **Hilb**. In presence of such a dagger one calls a map $f\colon X \to X$ positive, if it can be written of the form $f = g^\dagger \circ g$.
  In this context it makes sense to call a predicate $p = \langle p_1, p_2\rangle\colon X \to X \oplus X$ positive if both maps $p_i\colon X \to X$ are positive. But if $p_1 \geq 0$ and $p_2 = \mathrm{id} - p_1 \geq 0$, we must have $0 \leq p_1 \leq \mathrm{id}$. Such maps are called *effects*, see [45,43]. They are "quantum fuzzy predicates", and form one of the main examples of the notion of effect algebra. They are also called "unsharp" predicates, where the subset of projections contains the "sharp" predicates, see (the title of) [21]. These effects occur as predicates in the quantum weakest precondition calculus of [16]. It is noteworthy that the new representation (1) of positive predicates in Hilbert spaces captures effects, and not projections (the representations of formulas in the orthomodular logic started by von Neumann).

Since all these examples of predicates of the form $X \to X + X$ have the structure of an effect algebra, one immediately wonders if this structure exists in general for such predicates. It turns out that the coproduct has to satisfy certain assumptions, but if this is the case, we can define a partial sum operation $\ovee$ on predicates. This $\ovee$ is an essential part of the notion of an effect algebra. An example is the sum operation $+$ on the unit interval $[0, 1]$: it is clearly partial, since if $r + s > 1$ for $r, s \in [0, 1]$, their sum $r + s$ does not exist in $[0, 1]$. Interestingly, the founder of mathematical logic, George Boole, originally defined disjunction only as a partial operation, defined on disjoint subsets only. This makes perfect sense from the effect algebra perspective (see Example 7 below).

So assume we have two predicates $p, q\colon X \to X + X$ on X. We say that the sum $p \ovee q\colon X \to X + X$ exists, if there is a "bound" $b\colon X \to (X + X) + X$ such that:

$$\begin{tikzcd}[column sep=small]
 & (X \arrow[d, mapsto] & +\,X)+ \arrow[d, mapsto] & X \arrow[dl, mapsto] \\
X \arrow[ur, "b"] \arrow[r, "p"'] & X & +\,X &
\end{tikzcd}
\qquad
\begin{tikzcd}[column sep=small]
 & (X \arrow[dr, mapsto] & +\,X)+ \arrow[dl, mapsto] & X \arrow[dl, mapsto] \\
X \arrow[ur, "b"] \arrow[r, "q"'] & X & +\,X &
\end{tikzcd}
\qquad
\begin{tikzcd}[column sep=small]
 & (X \arrow[d, mapsto] & +\,X)+ \arrow[dl, mapsto] & X \arrow[dl, mapsto] \\
X \arrow[ur, "b"] \arrow[r, "p \ovee q"'] & X & +\,X &
\end{tikzcd}$$

Thus, the bound tries to combine the left parts of p and q in its first two components of $(X + X) + X$. For a more precise description, see Definition 3 below. In the three examples listed above, $\ovee$ is:

- union of subsets, but only if the subsets are disjoint;
- pointwise sum of fuzzy predicates $\varphi, \psi\colon X \to [0, 1]$, provided that $\varphi(x) + \psi(x) \leq 1$ for each $x \in X$;
- pointwise sum of effects, provided the sum is below the identity map.

This definition of sum $\ovee$ follows [35] where it is used only in a probabilistic context. In fact, the essence of many of the effect algebra results described below occur in [35], but in a more restricted setting.

In Section 6 it will be shown that for these predicates of the form $X \to X + X$ appropriate substitution functors f^* can be defined: for a map $f\colon X \to Y$, this f^* is a map of effect algebras from (positive) predicates on Y to (positive) predicates on X. The standard way to organise predicates and substitution in categorical logic is via indexed or fibred categories, see [32]. Such structures also arise in the current setting, with fibred effect structure, and even with a comprehension operation.

In the quantum case the substitution functor f^* takes the form:

$$f^*\Big(Y \xrightarrow{q} Y + Y\Big) = \Big(X \xrightarrow{f} Y \xrightarrow{q} Y + Y \xrightarrow{f^\dagger + f^\dagger} X + X\Big),$$

where f is a unitary (or more generally, a dagger monic, or isometric) map. This double occurrence of f, both before and after the predicate q is significant, since it gives rise to the Born rule. This rule says that the probability of a certain outcome is given by the square of the absolute value of the relevant amplitude. The double occurrence of f and $f^\dagger$ in substitution f^* leads directly to the Born rule (see Example 13), since $z^\dagger z = |z|^2$.

The final section concentrates on measurement in the present context. We view predicates as specifications of measurements—where in other settings projections, self-adjoints, or Positive Operator-Valued Measures (POVMs) are used, see [51]. It is first shown that in all our examples one can associate a certain characteristic map char_p with a predicate p, so that p can be recovered via substitution as $p = (\mathit{char}_p)^*(\Omega)$, from a special predicate Ω. This resembles the situation of a topos. The characteristic map char_p is used for two purposes: (1) to define test operations $\langle p?\rangle(-)$ and $[p?](-)$ like in dynamic logic, and (2) for the passage to successor states in measurement (specified by p). Measurement will be described in three different forms: on pure states, on predicate-transformers, and on mixed states (via duality). This works in our three flavours of logic: set-theoretic, probabilistic, and quantum, although in the first two cases measurement is a bit trivial, with $\mathit{char}_p = p$. In the quantum case we have $\mathit{char}_p = \sqrt{p}$, see Section 9 for more details. The square root acts as inverse for the squaring used in substitution (as mentioned above).

The novelty of this work lies first of all in the new way of representing predicates as maps $X \to X + X$, and in the description of the associated effect structure to capture the quantitative character of classical, probabilistic, and quantum logic. But on a deeper level three new elements can be distinguished.

(1) The Born rule provides the link between the mathematical formalism of quantum theory and experiments, and as such is almost single-handedly responsible for practically all predictions of quantum physics. Whereas Born in 1926 simply postulated this rule (in [8], see also [6]), attempts to derive the rule from first (physical) principles go back at least to the 1960s. The reference [44] contains an overview and discussion, and ends with the sentence: "The conclusion seems to be that no generally accepted derivation of the Born rule has been given to date, but this does not imply that such a derivation is impossible in principle.". Here the Born rule simply follows from the categorical organisation of effects in an indexed/fibred category. Of course, this does not provide a derivation of the Born rule from physical principles, but a characterisation in (categorical) logic, namely as substitution.

(2) Eigenvalues are frequently seen as irrelevant in quantum measurement, and are used only as labels or place holders. As illustration we quote [55, §§4.3.1]: "In particular, since the eigenvalues are simply labels for the subspaces or possible outcomes, the specific values of the eigenvalues are irrelevant; it matters only which ones are distinct." In the current setting, let $q\colon H \to H$ be a predicate/effect on a finite-dimensional Hilbert space H. In its spectral decomposition $q = \sum_i \lambda_i | v_i \rangle\langle v_i |$ the eigenvalues λ_i are in the unit interval $[0, 1]$. Hence, the combined eigenvalues form a fuzzy predicate $n \to [0, 1]$, where $n = \dim(H)$. These eigenvalues do play a role in the way we understand measurement here, namely as weights (multiplication factors) for the probabilities of the corresponding projection outcomes (onto the eigenvectors $| v_i \rangle$). The square roots $\sqrt{q}$, and hence also $\sqrt{\lambda_i}$, are essential for this to work. Analogously, in the probabilistic case, measurement wrt. a fuzzy predicate $\varphi\colon X \to [0, 1]$ means that the probability of $x \in X$ in a given state is multiplied by a factor $\varphi(x) \in [0, 1]$.

(3) A semantic foundation is given for a dynamic logic, involving modal operators $[f](p) = f^*(p)$ by substitution, and tests $\langle p?\rangle(q)$ and $[p?](q)$ defined via charac-

teristic maps *char*$_p$, see Subsection 9.1. The latter two operations have the familiar meaning in the set-theoretic and probabilistic case (including the Reichenbach implication), but seem to be new in the quantum case. Their use is illustrated in probability calculations in the familiar polarisation experiment (see Example 19).

The categorical study of quantum foundations started with the recognition of the sheaf-theoretic nature of the Kochen-Specker Theorem in [10][1], and with the use of tensor categories in [1] to capture composite systems and entanglement. The current work is in a sense orthogonal to the latter work. Here, the logical approach is not (yet) elaborated for composite systems, and in [1] the quantitative and probabilistic aspects of the theory are missing. Future work should lead to an appropriate combination.

This paper is organised as follows. First, it contains a discussion of coproducts (and biproducts), and especially of the conditions that they will have to satisfy in order to get the current approach off the ground. Section 3 describes the new approach to predicates in detail, with many examples, and Section 4 introduces the partial sum operation $\ovee$ on these predicates. Subsequently, Section 5 puts this operation $\ovee$ in the wider perspective of effect algebras/monoids/modules. The topic of substitution for predicates is covered in Section 6, and the effect monoid and module structure on predicates is defined in Sections 7 and 8. Substitution is used to define characteristic maps *char*$_p$ in Section 9, which give rise to test operations $[p?]$ and $\langle p?\rangle$. Finally, measurement, using these characteristic maps, appears in Section 10. The paper concludes with some final remarks and topics for further research.

2 Coproducts and biproducts

As explained in the introduction we will be using maps of the form $X \to X + X$ as predicates. This does not always make sense: for instance, in a poset (as a category) with join $\vee$ as coproduct $+$ one has $X \vee X = X$ so that predicates trivialise. In this section we will discuss some conditions on coproducts that will be relevant in the current setting. Special attention will be devoted to coproducts which are at the same time products (the so-called biproducts).

We start by recalling some basic facts about finite coproducts $(0, +)$ in an arbitrary category **C**. We write 0 for the initial object in **C**, with unique map $!\colon 0 \to X$ to each object $X \in$ **C**. The coproduct object $X + Y$ of two objects $X, Y \in$ **C** comes with two coprojections $\kappa_1 \colon X \to X + Y$ and $\kappa_2 \colon Y \to X + Y$ which are universal: for each pair of maps $f \colon X \to Z$ and $g \colon Y \to Z$ there is a unique cotuple $[f, g] \colon X + Y \to Z$ with $[f, g] \circ \kappa_1 = f$ and $[f, g] \circ \kappa_2 = g$. For two maps $h \colon X \to A$ and $k \colon Y \to B$ one writes $h + k = [\kappa_1 \circ h, \kappa_2 \circ k] \colon X + Y \to A + B$. The codiagonal map $X + X \to X$ is written as $\nabla = [\mathrm{id}, \mathrm{id}]$.

In the sequel we shall make frequent use of certain assumptions about coproducts $+$. At this stage we already make them explicit. These assumptions slightly generalise the ones used in [35].

---

[1] In the resulting topos-theoretic approach the goal is to interpret quantum mechanics within the left branch in Figure 1, see also [19,31,12].

Assumption 1 *(i). Specific diagram involving coproducts are pullbacks, namely:*

$$\begin{tikzcd}
A+X \arrow[r, "\mathrm{id}+g"] \arrow[d, "f+\mathrm{id}"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "\textbf{\textit{(E)}}"] & A+Y \arrow[d, "f+\mathrm{id}"] \\
B+X \arrow[r, "\mathrm{id}+g"'] & B+Y
\end{tikzcd}
\qquad
\begin{tikzcd}
X \arrow[r, "f"] \arrow[d, "\kappa_1"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "\textbf{\textit{(K)}}"] & Y \arrow[d, "\kappa_1"] \\
X+A \arrow[r, "f+\mathrm{id}"'] & Y+A
\end{tikzcd}
\qquad
\begin{tikzcd}
X \arrow[r, "f"] \arrow[d, "\kappa_1"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "\textbf{\textit{(K+)}}"] & Y \arrow[d, "\kappa_1"] \\
X+A \arrow[r, "f+g"'] & Y+B
\end{tikzcd}$$

The letter 'E' stands for exchange, 'K' for kappa (or coprojection).

(ii). The two parallel maps:

$$\begin{tikzcd}
(X+X)+X \arrow[r, bend left, "{[\mathrm{id},\kappa_2]}"] \arrow[r, bend right, "{[[\kappa_2,\kappa_1],\kappa_2]}"'] & X+X
\end{tikzcd}$$

acting as



(2)

are jointly monic; this means that for parallel maps f, g *one has* $f = g$ *as soon as both:*

$$[\mathrm{id},\kappa_2]\circ f = [\mathrm{id},\kappa_2]\circ g \quad \textit{and} \quad [[\kappa_2,\kappa_1],\kappa_2]\circ f = [[\kappa_2,\kappa_1],\kappa_2]\circ g.$$

(iii). Also the following diagrams are pullbacks,

$$\begin{tikzcd}
X+X \arrow[r, "f+f"] \arrow[d, "\nabla"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "\textbf{\textit{(D)}}"] & Y+Y \arrow[d, "\nabla"] \\
X \arrow[r, "f"'] & Y
\end{tikzcd}
\qquad
\begin{tikzcd}
(X+X)+X \arrow[r, "(f+f)+f"] \arrow[d, "\nabla+\mathrm{id}"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "\textbf{\textit{(D+)}}"] & (Y+Y)+Y \arrow[d, "\nabla+\mathrm{id}"] \\
X+X \arrow[r, "f+f"'] & Y+Y
\end{tikzcd}$$

where the letter 'D' stands for diagonal.

2.1 Extensive categories

We recall that coproducts are called *disjoint* if the coprojections κ_1, κ_2 are monic and their intersection is empty. The latter means that the following diagram is a pullback.

$$\begin{tikzcd}
0 \arrow[r, "!"] \arrow[d, "!"'] \arrow[dr, phantom, "\lrcorner", very near start] & Y \arrow[d, tail, "\kappa_2"] \\
X \arrow[r, tail, "\kappa_1"'] & X+Y
\end{tikzcd}$$

These coproducts are called *universal* if for each map $f\colon Z \to X+Y$ the two pullbacks below exist, and yield an isomorphism $[k_1,k_2]\colon Z_1+Z_2 \xrightarrow{\cong} Z$.

$$\begin{tikzcd}
Z_1 \arrow[r, tail, "k_1"] \arrow[d, "f_1"'] \arrow[dr, phantom, "\lrcorner", very near start] & Z \arrow[d, "f"] & Z_2 \arrow[l, tail, "k_2"'] \arrow[d, "f_2"] \arrow[dl, phantom, "\llcorner", very near start] \\
X \arrow[r, tail, "\kappa_1"'] & X+Y & Y \arrow[l, tail, "\kappa_2"]
\end{tikzcd}$$

A category with such disjoint and universal coproducts is called *extensive* [11]. For instance, the category **Sets**, like any topos (see *e.g* [7, Vol.3, §§5.9]), is extensive.

We remark that the initial object 0 of an extensive category is *strict*: any map $X \to 0$ is an isomorphism. In case an extensive category has products $\times$, these products distribute over coproducts, see [11].

Lemma 1. *In an extensive category Assumption 1 holds: all listed squares are pullbacks, and the two maps* $[\mathrm{id}, \kappa_2], [[\kappa_2, \kappa_1], \kappa_2]$ *are jointly monic.*

Proof. Elementary, but laborious. □

2.2 Biproducts in dagger categories

A *biproduct category* has biproducts $(0, \oplus)$; these are both products and coproducts, in the following manner. The object 0 is a *zero object*, which means that it is both initial and final. As a result, between any two objects X, Y there is a *zero map* $X \to 0 \to Y$, also written as 0. The biproduct $X_1 \oplus X_2$ comes with both coprojections and projections:

$$X_i \xrightarrow{\kappa_i} X_1 \oplus X_2 \xrightarrow{\pi_j} X_j \qquad \text{where} \qquad \pi_j \circ \kappa_i = \begin{cases} \mathrm{id} & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

In that case (co)projections can be expressed as (co)tuples, via:

$$\pi_1 = [\mathrm{id}, 0] \qquad \pi_2 = [0, \mathrm{id}] \qquad \kappa_1 = \langle \mathrm{id}, 0 \rangle \qquad \kappa_2 = \langle 0, \mathrm{id} \rangle.$$

One sees that the two coprojections are monomorphisms and their pullback is the zero object 0.

In presence of such biproducts $(0, \oplus)$ each homset of maps $X \to Y$ is a commutative monoid, via the zero map 0 and addition:

$$f + g = \Big(X \xrightarrow{\Delta = \langle \mathrm{id}, \mathrm{id} \rangle} X \oplus X \xrightarrow{f \oplus g} Y \oplus Y \xrightarrow{\nabla = [\mathrm{id}, \mathrm{id}]} Y\Big). \tag{3}$$

This monoid structure is preserved by pre- and post-composition, making the category enriched over the category **CMon** of commutative monoids. Moreover, (co)tupling and $+$ interact in the following way.

$$\begin{array}{rclcrcl} [f, g] & = & (f \circ \pi_1) + (g \circ \pi_2) & \qquad & \langle f_1, f_2 \rangle + \langle g_1, g_2 \rangle & = & \langle f_1 + g_1, f_2 + g_2 \rangle \\ \langle f, g \rangle & = & (\kappa_1 \circ f) + (\kappa_2 \circ g) & \qquad & [f_1, f_2] + [g_1, g_2] & = & [f_1 + g_1, f_2 + g_2] \end{array}$$

In an *additive* category these commutative monoids have additive inverses $-f \colon X \to Y$, for each $f \colon X \to Y$, making the category **Ab**-enriched (where **Ab** is the category of Abelian groups).

A category **C** is a *dagger category* if it comes with a functor $(-)^\dagger \colon \mathbf{C}^{\mathrm{op}} \to \mathbf{C}$ that is the identity on objects; on arrows it should satisfy $f^{\dagger\dagger} = f$. In such a dagger category a map f with $f^\dagger \circ f = \mathrm{id}$ is called a *dagger mono* (or isometry); if $f \circ f^\dagger = \mathrm{id}$ one calls f a *dagger epi*. A map that is both a dagger mono and a dagger epi is called a *dagger* *iso*, or a *unitary* map. We write $\dagger$-*Iso*$(\mathbf{C}) \hookrightarrow \dagger$-*Mono*$(\mathbf{C}) \hookrightarrow \mathbf{C}$ for the subcategories with all objects from **C**, and unitary / dagger monic maps only. Finally, an endomap $f\colon X \to X$ is called *positive* if it can be written as $f = g^\dagger \circ g$, for some g.

A *dagger biproduct* category is a dagger category with biproducts in which $(\pi_i)^\dagger = \kappa_i$ — and thus $(\kappa_i)^\dagger = \pi_i$ and $\langle f, g\rangle^\dagger = [f^\dagger, g^\dagger]$ and $[f, g]^\dagger = \langle f^\dagger, g^\dagger\rangle$ and $(f \oplus g)^\dagger = f^\dagger \oplus g^\dagger$. It is easy to see that if $f, g\colon X \to X$ are both positive, then so is the sum $f + g$ from (3). Such a dagger biproduct category is called *zerosumfree* if $f + g = 0 \Rightarrow f = g = 0$, for positive maps f, g.

A *dagger additive* category is a dagger biproduct category with additive inverses in homsets.

Lemma 2. *With respect to Assumption 1,*

(i). in a biproduct category, diagrams (E) and (K) are pullbacks, and diagram (K+) too when the map g is zero monic (that is $g \circ h = 0 \Rightarrow h = 0$);

(ii). in an additive category, the two maps $(X \oplus X) \oplus X \rightrightarrows X \oplus X$,

$$[\mathrm{id}, \kappa_2] = \langle \pi_1 \circ \pi_1, (\pi_2 \circ \pi_1) + \pi_2\rangle \qquad [[\kappa_2, \kappa_1], \kappa_2] = \langle \pi_2 \circ \pi_1, (\pi_1 \circ \pi_1) + \pi_2\rangle$$

from Assumption 1.(ii) are jointly monic;

(iii). in a dagger biproduct category, diagrams (D) and (D+) are pullbacks if the map f is unitary.

Proof. (i). In order to check that the square (E) in Assumption 1.(i) is a pullback, assume $h = \langle h_1, h_2\rangle\colon Z \to B \oplus X$ and $k = \langle k_1, k_2\rangle\colon Z \to A \oplus Y$ satisfy $(\mathrm{id} \oplus g) \circ h = (f \oplus \mathrm{id}) \circ k$. This means $h_1 = f \circ k_1$ and $g \circ h_2 = k_2$. Hence $\ell = \langle k_1, h_2\rangle$ is the (unique) mediating map:

$$\begin{aligned}(f \oplus \mathrm{id}) \circ \ell &= \langle f \circ k_1, h_2\rangle \\ &= \langle h_1, h_2\rangle = h\end{aligned} \qquad \begin{aligned}(\mathrm{id} \oplus g) \circ \ell &= \langle k_1, g \circ h_2\rangle \\ &= \langle k_1, k_2\rangle = k.\end{aligned}$$

For diagram (K+), assume we have $h = \langle h_1, h_2\rangle\colon Z \to X \oplus A$ and $k\colon Z \to Y$ with $(f \oplus g) \circ h = \kappa_1 \circ k$. Then $\langle f \circ h_1, g \circ h_2\rangle = \langle k, 0\rangle$. Hence $f \circ h_1 = k$ and $g \circ h_2 = 0$. Since g is assumed to be zero monic, we get $h_2 = 0$. Thus we have $h_1\colon Z \to X$ as unique mediating map.

(ii). Assume $f = \langle\langle f_1, f_2\rangle, f_3\rangle\colon A \to (X \oplus X) \oplus X$ and $g = \langle\langle g_1, g_2\rangle, g_3\rangle\colon A \to (X \oplus X) \oplus X$ satisfy $[\mathrm{id}, \kappa_2] \circ f = [\mathrm{id}, \kappa_2] \circ g$ and $[[\kappa_2, \kappa_1], \kappa_2] \circ f = [[\kappa_2, \kappa_1], \kappa_2] \circ g$. This means $\langle f_1, f_2 + f_3\rangle = \langle g_1, g_2 + g_3\rangle$ and $\langle f_2, f_1 + f_3\rangle = \langle g_2, g_1 + g_3\rangle$. Hence:

$$f_1 = g_1 \qquad f_2 + f_3 = g_2 + g_3 \qquad f_2 = g_2 \qquad f_1 + f_3 = g_1 + g_3.$$

Then: $f_3 = (g_1 + g_3) - f_1 = (g_1 + g_3) - g_1 = g_3$. Hence $f = g$.

(iii). Easy. $\square$

Our main examples of dagger additive categories are the categories **FdHilb** and **Hilb** of finite-dimensional and arbitrary Hilbert spaces. The morphisms in these categories are the linear continuous (equivalently, bounded) maps. These categories also have a (symmetric) monoidal structure $(\mathbb{C}, \otimes)$, where $\otimes$ distributes over biproducts $\oplus$. The additive monoid of positive operators $H \to H$ on a Hilbert space H is zerosumfree: $A + B = 0 \Longrightarrow A = B = 0$.[2]

Another example is the category **BifMRel** of sets and bifinite multirelations (used for instance in [34]). Objects in **BifMRel** are sets, and morphisms $X \to Y$ are complex-valued relations $r\colon X \times Y \to \mathbb{C}$, with a "bifiniteness" requirement: for each $x \in X$ and $y \in Y$ one has finite sets:

$$\{v \in Y \mid r(x, v) \neq 0\} \qquad \text{and} \qquad \{u \in X \mid r(u, y) \neq 0\}.$$

The identity map $X \to X$ in **BifMRel** is the identity relation $\mathrm{id}_X\colon X \times X \to \mathbb{C}$, given by $\mathrm{id}_X(x, x) = 1$, and $\mathrm{id}_X(x, x') = 0$ when $x \neq x'$. The composition $s \circ r$ in **BifMRel**, for $r\colon X \to Y$ and $s\colon Y \to Z$ is the relational composition $X \times Z \to \mathbb{C}$ given by:

$$(s \circ r)(x, z) \,=\, \textstyle\sum_{y \in Y} r(x, y) \cdot s(y, z).$$

It is not hard to see that **BifMRel** is a dagger category, with dagger functor $(-)^\dagger$ given as identity on objects and complex conjugation on morphisms. Thus $r\colon X \to Y$ yields $r^\dagger\colon Y \to X$ given by $r^\dagger(y, x) = \overline{r(x, y)}$.

The category **BifMRel** also has (dagger) biproducts, given by coproducts $(0, +)$ on the underlying sets. Products $(1, \times)$ on these sets yield a (dagger) symmetric monoidal structure on **BifMRel**, that distributes over the biproducts.

The category **BifMRel** allows relatively easy descriptions of (discrete) quantum systems, see *e.g.* [34] for quantum walks in **BifMRel**. Also, a quantum Turing machine with state space S and set of symbols Σ may be represented as dagger mono $r\colon S \times \Sigma \to (S \times \Sigma) + (S \times \Sigma)$ in **BifMRel**, where the biproduct $+$ is used to describe the two possible successor options, namely tape head moving left or right (see also *e.g.* [58]). The requirement that r be a dagger mono ensures that for each $(s, a) \in S \times \Sigma$ one has $\sum_z |r((s, \sigma), z)|^2 = 1$, that is, the amplitudes associated with transitions to successors of (s, a) satisfy the usual requirement to obtain a quantum state.

2.3 Monads and Kleisli categories

In this subsection we briefly review three well-known monads on **Sets**, namely the distribution monad $\mathcal{D}$, the multiset monad $\mathcal{M}$ and the powerset monad $\mathcal{P}$. We shall review Assumption 1, especially for $\mathcal{D}$ and $\mathcal{P}$.

We assume the reader is familiar with the notion of monad (see [4,48]). For a monad $T = (T, \eta, \mu)$ on a category **C** we write $\mathcal{K}\ell(T)$ for the Kleisli category of T. Its objects are objects $X \in$ **C** and its morphisms $f\colon X \to Y$ are maps $f\colon X \to T(X)$ in **C**. For such a map we write $f^\# = \mu \circ T(f)\colon T(X) \to T(Y)$. Composition $f \circ g$ in $\mathcal{K}\ell(T)$ is then given by $f^\# \circ g$ in **C**; the identity $X \to X$ in $\mathcal{K}\ell(T)$ is the unit $\eta\colon X \to T(X)$. There is a standard functor $\mathcal{J}\colon$ **C** $\to \mathcal{K}\ell(T)$, which is the identity on objects and maps $f\colon X \to Y$ to $\mathcal{J}(f) = \eta \circ f\colon X \to T(Y)$. If **C** has coproducts $(0, +)$, then so has

---

[2] The proof uses: if a (non-negative, real-valued) inner product $\langle x \,|\, Ax \rangle$ is 0, for all $x \in H$, implies $A = 0$, essentially because a bilinear form is determined by its diagonal values (see *e.g.* [27, §§I.3] or [29, Proposition 1.21]).

$\mathcal{K}\ell(T)$, with coprojections $\mathcal{J}(\kappa_i)$ and tupling of Kleisli maps f, g given by tupling $[f, g]$ as in **C**. In all our examples **C** $=$ **Sets**, so Kleisli categories of the monads we consider have coproducts.

We turn to the multiset monad $\mathcal{M}$ on **Sets**. A multiset is like a subset except that elements may occur multiple times. Hence one needs a way of counting elements. Most generally this can be done in a semiring, but in the current setting we count in the complex numbers $\mathbb{C}$. Thus the collection of (complex-valued, finite) multisets of a set X is defined in terms of formal linear combinations of elements of X, as in:

$$\mathcal{M}(X) = \{z_1x_1 + \cdots + z_nx_n \mid z_i \in \mathbb{C} \text{ and } x_i \in X\}. \tag{4}$$

Such a formal sum $\sum_i z_ix_i$ can also be described as a function $m\colon X \to \mathbb{C}$ with finite support, that is, where *supp*$(m) = \{x \mid m(x) \neq 0\}$ is finite.

This mapping $X \mapsto \mathcal{M}(X)$ is functorial: for a function $f\colon X \to Y$ one defines $\mathcal{M}(f)\colon \mathcal{M}(X) \to \mathcal{M}(Y)$ by $\mathcal{M}(f)(\sum_i z_ix_i) = \sum_i z_if(x_i)$. There is a unit $\eta\colon X \to \mathcal{M}(X)$ given by $\eta(x) = 1x$, and a multiplication $\mu\colon \mathcal{M}^2(X) \to \mathcal{M}(X)$ described by:

$$\mu(\textstyle\sum_i z_im_i) = \sum_{i,j}(z_i \cdot w_{ij})x_{ij} \qquad \text{where} \qquad m_i = \sum_j w_{ij}x_{ij}.$$

It is not hard to see that $\mathcal{M}(X)$ is the free vector space (over $\mathbb{C}$) on the set X.

Remark 1. We shall not use the multiset monad $\mathcal{M}$ explicitly, but it is used implicitly in the category **BifMRel** of sets and bifinite multirelations. The bifiniteness requirement for a map $r\colon X \to Y$ in **BifMRel** means that r factors both as $X \to \mathcal{M}(Y)$ and as $Y \to \mathcal{M}(X)$. Thus, by freeness, r gives rise to two linear maps $\mathcal{M}(X) \to \mathcal{M}(Y)$ and $\mathcal{M}(Y) \to \mathcal{M}(X)$ in opposite directions. In this category **BifRel** the objects/sets X can be seen as the computational basis, with associated free vector space $\mathcal{M}(X)$.

We turn to the discrete probability distribution monad $\mathcal{D}$. It is a more restricted version of the multiset monad that sends a set to formal convex combinations of its elements:

$$\mathcal{D}(X) = \{r_1x_1 + \cdots + r_nx_n \mid r_i \in [0,1] \text{ and } x_i \in X \text{ and } \textstyle\sum_i r_i = 1\}. \tag{5}$$

Elements of $\mathcal{D}(X)$ are functions $m\colon X \to [0,1]$ with finite support and $\sum_x m(x) = 1$.

The action of $\mathcal{D}$ on functions is essentially the same as $\mathcal{M}$; also $\mathcal{M}$'s unit and multiplication restrict to $\mathcal{D}$. In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the distribution monad a map $X \to Y$ is a function $f\colon X \to \mathcal{D}(Y)$. It may also be described as a stochastic matrix $M_f\colon X \times Y \to [0,1]$, where $M_f(x,y) = f(x)(y)$. For each $x \in X$ the subset $\{y \in Y \mid M_f(x,y) \neq 0\}$ is finite, with $\sum_y M_f(x,y) = 1$. Thus, such a matrix M_f corresponds to a stochastic matrix.

Lemma 3. *In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the distribution monad, the diagrams (E) and (K+)—and thus (K)—from Assumption 1.(i) are pullbacks, and the two maps in Assumption 1.(ii) are jointly monic.*

Proof. We check that diagram (E) is a pullback and leave (K+) to the reader. We reason pointwise and assume $\varphi \in \mathcal{D}(B+X)$ and $\psi \in \mathcal{D}(A+Y)$ satisfy $(\mathrm{id}+g)^{\#}(\varphi) = (f+\mathrm{id})^{\#}(\psi)$. Thus we have equal functions $B+Y \to [0,1]$. This means for each $b \in B$ and $y \in Y$,

$$\varphi(\kappa_1 b) \;=\; \textstyle\sum_{a\in A} \psi(\kappa_1 a)\cdot f(a)(b) \qquad\qquad \textstyle\sum_{x\in X} \varphi(\kappa_2 x)\cdot g(x)(y) \;=\; \psi(\kappa_2 y)$$

Hence we define $\chi \in \mathcal{D}(A+X)$ as $\chi(\kappa_1 a) = \psi(\kappa_1 a)$ and $\chi(\kappa_2 x) = \varphi(\kappa_2 x)$. This yields a probability distribution:

$$\begin{aligned}
\textstyle\sum_{u\in A+X} \chi(u) &= \textstyle\sum_a \chi(\kappa_1 a) \;+\; \sum_x \chi(\kappa_2 x) \\
&= \textstyle\sum_a \psi(\kappa_1 a) \;+\; \sum_x \varphi(\kappa_2 x) \\
&= \textstyle\sum_a \psi(\kappa_1 a)\cdot(\sum_b f(a)(b)) \;+\; \sum_x \varphi(\kappa_2 x) \\
&= \textstyle\sum_{a,b} \psi(\kappa_1 a)\cdot f(a)(b) \;+\; \sum_x \varphi(\kappa_2 x) \\
&= \textstyle\sum_b \varphi(\kappa_1 b) \;+\; \sum_x \varphi(\kappa_2 x) \\
&= 1.
\end{aligned}$$

Also, this χ is the appropriate mediating distribution:

$$\begin{aligned}
(f+\mathrm{id})^{\#}(\chi)(\kappa_1 b) &= \textstyle\sum_{a\in A} \chi(\kappa_1 a)\cdot f(a)(b) \\
&= \textstyle\sum_{a\in A} \psi(\kappa_1 a)\cdot f(a)(b) \;=\; \varphi(\kappa_1 b) \\
(f+\mathrm{id})^{\#}(\chi)(\kappa_2 x) &= \chi(\kappa_2 x) \;=\; \varphi(\kappa_2 x) \\
(\mathrm{id}+g)^{\#}(\chi)(\kappa_1 a) &= \chi(\kappa_1 a) \;=\; \psi(\kappa_1 a) \\
(\mathrm{id}+g)^{\#}(\chi)(\kappa_2 y) &= \textstyle\sum_x \chi(\kappa_2 x)\cdot g(x)(y) \\
&= \textstyle\sum_x \varphi(\kappa_2 x)\cdot g(x)(y) \;=\; \psi(\kappa_2 y).
\end{aligned}$$

Thus, $(f+\mathrm{id})^{\#}(\chi) = \varphi$ and $(\mathrm{id}+g)^{\#}(\chi) = \psi$. Uniqueness is left to the reader.

Finally, we check that the two maps $[\mathrm{id}, \kappa_2], [[\kappa_2, \kappa_1], \kappa_2] \colon (X+X)+X \to X+X$ are jointly monic in $\mathcal{K}\ell(\mathcal{D})$. So assume $\varphi, \psi \in \mathcal{D}((X+X)+X)$ satisfy:

$$\mathcal{D}([\mathrm{id},\kappa_2])(\varphi) \;=\; \mathcal{D}([\mathrm{id},\kappa_2])(\psi) \qquad\qquad \mathcal{D}([[\kappa_2,\kappa_1],\kappa_2])(\varphi) \;=\; \mathcal{D}([[\kappa_2,\kappa_1],\kappa_2])(\psi).$$

This means for each $x \in X$,

$$\begin{array}{ll}
\varphi(\kappa_1\kappa_1 x) \;=\; \psi(\kappa_1\kappa_1 x) \qquad & \varphi(\kappa_1\kappa_2 x) + \varphi(\kappa_2 x) \;=\; \psi(\kappa_1\kappa_2 x) + \psi(\kappa_2 x) \\
\varphi(\kappa_1\kappa_2 x) \;=\; \psi(\kappa_1\kappa_2 x) \qquad & \varphi(\kappa_1\kappa_1 x) + \varphi(\kappa_2 x) \;=\; \psi(\kappa_1\kappa_1 x) + \psi(\kappa_2 x)
\end{array}$$

From this it follows that $\varphi(\kappa_2 x) = \psi(\kappa_2 x)$, and thus $\varphi = \psi$. □

Remark 2. The Kleisli category $\mathcal{K}\ell(\mathcal{P})$ of the powerset monad $\mathcal{P}$ can be identified with the category of sets and relations between them. It does not satisfy Assumption 1.(ii). We construct a counterexample to the “jointly monic” requirement.

For a singleton set 1 we have $(1+1)+1 \cong \{1,2,3\}$ and $1+1 \cong \{1,2\}$. Writing $p = [\mathrm{id}, \kappa_2]$ and $q = [[\kappa_2, \kappa_1], \kappa_2]$ in Assumption 1.(ii), we can describe $p, q \colon 3 \rightrightarrows 2$ as:

$$p(1) = 1,\ p(2) = p(3) = 2 \qquad\qquad q(1) = q(3) = 2,\ q(2) = 1.$$

The two subset $U = \{1,2\}$ and $V = \{1,2,3\}$ of 3 are clearly different but they satisfy:

$$\mathcal{P}(p)(U) \;=\; \{1,2\} \;=\; \mathcal{P}(p)(V) \qquad \text{and} \qquad \mathcal{P}(q)(U) \;=\; \{1,2\} \;=\; \mathcal{P}(q)(V).$$

3 Predicates

In this section we introduce the new representation of predicates and describe some examples.

Definition 1. *Let* **C** *be a category with coproducts* $+$. *For each object* $X \in$ **C** *we define a collection* Pred(X) *of predicates on* X *as:*

$$\mathit{Pred}(X) \;=\; \{p \colon X \to X + X \mid \nabla \circ p = \mathrm{id}_X\},$$

where $\nabla = [\mathrm{id}, \mathrm{id}] \colon X + X \to X$ *is the codiagonal.*

In case **C** *is a dagger biproduct category, a predicate* $p \colon X \to X \oplus X$ *is called* positive *if both the components* $\pi_1 \circ p, \pi_2 \circ p \colon X \to X$ *are positive maps. We write* pPred(X) *for the collection of such positive predicates.*

There is a "truth" predicate $1 = \kappa_1 \colon X \to X + X$ and also a "false" predicate $0 = \kappa_2 \colon X \to X + X$. Of course, it is a matter of convention/choice which $+$ component one uses for truth or false. For each predicate $p \colon X \to X + X$ there is an orthocomplement $p^{\perp} = [\kappa_2, \kappa_1] \circ p$, obtained by swapping the $+$ components. Clearly, $p^{\perp\perp} = p$ and $0^{\perp} = 1$, and thus $1^{\perp} = 0$.

Definition 2. *Let our category* **C** *have besides coproduct* $+$ *also a monoidal structure* $(I, \otimes)$. *The (positive) predicates* Pred(I) *or* pPred(I) *on the tensor unit* I *will be called* probabilities. *We often write* Pr(**C**) *for this collection of probabilities in the category* **C**.

Later on, in Theorem 2, we shall see that probabilities carry a commutative monoid structure.

Before considering further structure on these predicates we illustrate what they are in several different categories. As we shall see in a dagger category the *positive* predicates are the interesting ones. Hence we sometimes drop the adjective 'positive' in such a setting.

Example 1. In the category **Sets** of sets and functions a predicate can be identified with a subset. Concretely, $U \subseteq X$ corresponds to $p_U \colon X \to X + X$ with $\nabla \circ p_U = \mathrm{id}$ via:

$$p_U(x) \;=\; \begin{cases} \kappa_1 x & \text{if } x \in U \\ \kappa_2 x & \text{if } x \notin U. \end{cases}$$

Thus, a predicate yields a partition of the set X into two disjoint subsets $U, \neg U$ that cover X. In this case *Pred*(X) is the set $\mathcal{P}(X)$ of subsets of X, with truth $1 = (X \subseteq X)$, false $0 = (\emptyset \subseteq X)$ and orthocomplement via negation: $(p_U)^{\perp} = p_{\neg u}$. The predicates on a singleton set 1 correspond to Booleans: *Pred*$(1) \cong \mathcal{P}(1) \cong \{0, 1\}$.

One other way to look at these predicates $X \to X + X$ in **Sets** is to use the isomorphisms:

$$X + X \;\cong\; (1 \times X) + (1 \times X) \;\cong\; (1 + 1) \times X \;\cong\; 2 \times X.$$

When we rephrase the condition $\nabla \circ p = \mathrm{id}$ via this isomorphism we see that a predicate corresponds a pair of maps $\langle \varphi, \mathrm{id} \rangle \colon X \to 2 \times X$, and thus to a "characteristic" function $\varphi \colon X \to 2$. This can be done in any distributive category. Predicates of this form $X \to 2$ have been studied in such categories for instance in [14]. Here we prefer to use the representation $X \to X + X$ because it highlights the similarities between predicates in different categories.

Example 2. Recall that the category of sets and relations can be identified with the Kleisli category $\mathcal{K}\ell(\mathcal{P})$ of the powerset monad $\mathcal{P}$. A predicate on a set X in this category is a map $p \colon X \to \mathcal{P}(X + X)$ with $\mathcal{P}(\nabla) \circ p = \{-\}$. This means that $p(x) \subseteq \{\kappa_1(x), \kappa_2(x)\}$ is a non-empty subset. Hence there are three options:

$$p(x) = \{\kappa_1 x\} \quad \text{ or } \quad p(x) = \{\kappa_2 x\} \quad \text{ or } \quad p(x) = \{\kappa_1 x, \kappa_2 x\}.$$

With such a predicate one naturally associates two subsets $U_1, U_2 \subseteq X$ given by:

$$U_1 = \{x \mid \kappa_1 x \in p(x)\} \qquad U_2 = \{x \mid \kappa_2 x \in p(x)\}.$$

Conversely, with such a pair of subsets U_1, U_2 one associates a predicate:

$$p_{U_1,U_2}(x) = \begin{cases} \{\kappa_1 x\} & \text{if } x \in U_1 - U_2 \\ \{\kappa_2 x\} & \text{if } x \in U_2 - U_1 \\ \{\kappa_1 x, \kappa_2 x\} & \text{if } x \in U_1 \cap U_2. \end{cases}$$

In this subset *Pred*$(X) \hookrightarrow \mathcal{P}(X) \times \mathcal{P}(X)$ of predicates on $X \in \mathcal{K}\ell(\mathcal{P})$ we have true $1 = (X, \emptyset)$ and false $0 = (\emptyset, X)$. Orthocomplement involves swapping: $(U_1, U_2)^{\perp} = (U_2, U_1)$.

The Kleisli category $\mathcal{K}\ell(\mathcal{P})$ of sets and relations is a dagger category (via relation reversal) and has biproducts $+$. Hence a predicate $p_{U_1,U_2} \colon X \to \mathcal{P}(X + X)$ corresponds to two maps $p_{U_1}, p_{U_2} \colon X \to \mathcal{P}(X)$, namely:

$$p_{U_i}(x) = \begin{cases} \{x\} & \text{if } x \in U_i \\ \emptyset & \text{otherwise.} \end{cases}$$

These maps p_{U_i} are projections: they satisfy $(p_{U_i})^{\dagger} = p_{U_i} = p_{U_i} \circ p_{U_i}$. Hence they are positive. Thus, a predicate in $\mathcal{K}\ell(\mathcal{P})$ is automatically positive.

Example 3. In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the distribution monad $\mathcal{D}$ a predicate p on a set X corresponds to a fuzzy predicate $\varphi\colon X \to [0,1]$. For such a predicate φ a Kleisli map $p_\varphi\colon X \to \mathcal{D}(X+X)$ with $\mathcal{D}(\nabla)(p(x)) = 1x$ is given by convex sums of the form:

$$p_\varphi(x) \,=\, \varphi(x)\kappa_1 x + (1-\varphi(x))\kappa_2 x.$$

In this set of fuzzy predicates we have a truth predicate $X \to [0,1]$ that is always 1, and a false predicate that is always 0. Orthocomplement is given by $\varphi^\perp(x) = 1 - \varphi(x)$.

The predicates on the singleton (final) set 1 can be identified with probabilities: *Pred*$(1) \cong [0,1]$.

Example 4. In the category **BifMRel** of sets and bifinite multirelations a predicate $p\colon X \to X + X$ corresponds to a bifinite map $p\colon X \times (X+X) \to \mathbb{C}$. The condition $\nabla \circ p = \mathrm{id}$ means, for $x, x' \in X$ with $x \neq x'$,

$$p(x,\kappa_1 x) + p(x,\kappa_2 x) \,=\, 1 \qquad \text{and} \qquad p(x,\kappa_1 x') + p(x,\kappa_2 x') \,=\, 0.$$

Thus, the 'κ_2-part' is determined by the 'κ_1-part', via:

$$p(x,\kappa_2 x') \,=\, \begin{cases} 1 - p(x,\kappa_1 x) & \text{if } x = x' \\ -p(x,\kappa_1 x') & \text{otherwise.} \end{cases}$$

Hence we may identify p with a matrix in $\mathbb{C}^{X\times X}$, via $(x,x') \mapsto p(x,\kappa_1 x')$, with only finitely many entries in each row and column.

A positive predicate p has probabilities $p(x,\kappa_1 x) \in [0,1]$ as diagonal values. In comparison with the fuzzy predicates in $[0,1]^X$ in the probabilistic world (see Example 3), the predicates in the quantum world given by **BifMRel** involve such a fuzzy predicate on the diagonal, but additionally involve other values in $\mathbb{C}$ on the non-diagonal entries. These other values may lead to the interferences that are typical of quantum computation — and are missing in a probabilistic setting.

The positive predicates on the tensor unit $1 \in$ **BifMRel** are the probabilities $[0,1]$.

Example 5. In the category **Vect** of vector spaces (over $\mathbb{C}$), the coproduct is at the same time a product, and thus a biproduct, written as $\oplus$. A map $p\colon V \to V \oplus V$ with $\nabla \circ p = \mathrm{id}$ thus consists of a pair of maps $p = \langle p_1, p_2\rangle$ with $p_1 + p_2 = \mathrm{id}$. Hence, the predicate p is determined by its first component p_1, since $p_2 = \mathrm{id} - p_1$.

The category **Hilb** of Hilbert spaces and linear continuous (bounded) maps is a dagger biproduct category. The positive predicates $\langle q, \mathrm{id} - q\rangle\colon H \to H \oplus H$ are given by a positive map $q\colon H \to H$ with $\mathrm{id} - q$ also positive. This means $0 \leq q \leq \mathrm{id}$. Such a map q is called an *effect* [20]. These effects are used as predicates for instance in the quantum weakest precondition calculus of [16], and in many other places.

Now suppose H has finite dimension, say $n \in \mathbb{N}$. For an effect $q\colon H \to H$ we can compute the eigenvalues $\lambda_1, \ldots, \lambda_n$, say with corresponding eigenvectors $e_i \in H$. The eigenvalues are real, non-negative numbers since q is positive. The eigenvectors form an orthonormal basis of H. Then:

$$\lambda_i \,=\, \lambda_i\langle e_i \,|\, e_i\rangle \,=\, \langle \lambda_i e_i \,|\, e_i\rangle \,=\, \langle q\, e_i \,|\, e_i\rangle \,\leq\, \langle \mathrm{id} e_i \,|\, e_i\rangle \,=\, 1.$$

Hence the eigenvalues $\lambda_i \in [0,1]$ are probabilities. We can put the eigenvectors as columns in a unitary matrix T so that $q = TDT^{\dagger}$, where D is a diagonal matrix with eigenvalues λ_i on the diagonal. These eigenvalues form a fuzzy predicate $\lambda \in [0,1]^n$, like in Example 3. Thus, a predicate q on a finite-dimensional Hilbert space is, up-to unitary change-of-base, a fuzzy predicate.

Of special interest are predicates on the complex numbers $\mathbb{C}$, considered as Hilbert space. This space $\mathbb{C}$ is the unit I for a tensor product $\otimes$ on **Hilb**. Linear (continuous) maps $\mathbb{C} \to \mathbb{C}$ can be identified with elements $z \in \mathbb{C}$. Thus, predicates on $I = \mathbb{C}$ correspond to complex numbers: *Pred*$(I) \cong \mathbb{C}$, since a predicate can be identified with a pair $\langle z, 1-z\rangle$ where $z \in \mathbb{C}$; *positive* predicates can be described as the unit interval: *pPred*$(I) \cong [0,1]$. Thus, the probabilities in **Hilb** are the ordinary probabilities.

One sees that in all of these illustrations, except Example 2 involving relations, the "left part" of a predicate determines the "right part". We shall see that the kind of predicates considered here does indeed not work well for the category of relations.

In the beginning of this section we have seen that κ_1 (true) and κ_2 (false) always exist as predicates $X \to X + X$. In the next example we describe a similar predicate that is always there. It will play a role in the description of characteristic maps, see Section 9.

Example 6. In a category with coproducts we have for each object X a map:

$$\mathit{\Omega}_X = \Big(X + X \xrightarrow{\kappa_1 + \kappa_2} (X+X) + (X+X) \Big).$$

This $\mathit{\Omega}_X$ is a predicate since:

$$\nabla \circ \mathit{\Omega}_X \;=\; \nabla \circ (\kappa_1 + \kappa_2) \;=\; [\kappa_1, \kappa_2] \;=\; \mathrm{id}.$$

The predicate $\mathit{\Omega}$ may be understood as "left", which is our choice for "truth": in the category **Sets** this $\mathit{\Omega}$ corresponds to the "left" subset $\kappa_1 \colon X \rightarrowtail X + X$. In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ it corresponds to the "left" fuzzy predicate $\psi \in [0,1]^{X+X}$ with $\psi(\kappa_1 x) = 1$ and $\psi(\kappa_2 x) = 0$, for each $x \in X$.

In a dagger biproduct category one has $\pi_i \circ \mathit{\Omega} = \kappa_i \circ \pi_i = (\pi_i)^{\dagger} \circ \pi_i$, so that $\mathit{\Omega}$ is automatically a positive predicate. If we consider $\mathit{\Omega}$ in the category **Hilb** of Hilbert spaces on the object $X = \mathbb{C}$ of complex numbers we may describe it as matrix:

$$\mathbb{C}^2 = \mathbb{C} \oplus \mathbb{C} \xrightarrow{\mathit{\Omega} = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{smallmatrix}\right)} \big(\mathbb{C} \oplus \mathbb{C}\big) \oplus \big(\mathbb{C} \oplus \mathbb{C}\big) = \mathbb{C}^4.$$

We conclude this section by briefly showing how the categorical representation of "sharp" quantum predicates as dagger kernels fits in the current setting.

Remark 3. In [30] dagger categories with dagger kernels are studied. Such a dagger kernel k is at the same time a kernel $k = \ker(f)$ of some map f, and a dagger mono (meaning: $k^{\dagger} \circ k = \mathrm{id}$). These dagger kernels with codomain X form an orthomodular lattice of predicates on X. The orthocomplement $k^\perp$ is defined as the kernel $\ker(k^\dagger)$, see [30] for details.

If one assumes that the dagger kernel category at hand also has dagger biproducts $\oplus$, then one can translate dagger kernels into positive predicates (as used here). This works as follows. Given a dagger kernel $k\colon U \rightarrowtail X$, the composite $k \circ k^\dagger\colon X \to X$ is a positive map. In a similar way $k^\perp \circ (k^\perp)^\dagger\colon X \to X$ is positive. Together they form a tuple:

$$p_k = \Big(X \xrightarrow{\langle k\circ k^\dagger, k^\perp\circ(k^\perp)^\dagger\rangle} X \oplus X\Big)$$

We claim that this p_k is a (positive) predicate on X, so we need to show $\nabla \circ p_k = \mathrm{id}$.

Let us write $U^\perp$ for the domain of the orthocomplement $k^\perp$. We claim that the following cotuple is an isomorphism:

$$U \oplus U^\perp \xrightarrow{[k,k^\perp]} X$$

The first thing to note is that this map is a dagger mono: the equation $[k,k^\perp]^\dagger \circ [k,k^\perp] = \mathrm{id}$ follows from an easy computation.

We shall use the basic result that the kernel of a tuple $\langle f, g\rangle$ is the intersection of kernels: $\ker(f) \wedge \ker(g)$. Then:

$$\begin{aligned} 0 \;=\; k^\perp \wedge k \;=\; k^\perp \wedge k^{\perp\perp} \;&=\; \ker(k^\dagger) \wedge \ker((k^\perp)^\dagger) \\ &=\; \ker(\langle k^\dagger, (k^\perp)^\dagger\rangle) \;=\; \ker([k,k^\perp]^\dagger) \;=\; [k,k^\perp]^\perp. \end{aligned}$$

We thus get $[k,k^\perp] = 1$, as subobject. We already saw that $[k,k^\perp]$ is a dagger mono. Hence it is an isomorphism, with $[k,k^\perp]^\dagger$ also as left inverse:

$$\begin{aligned} \mathrm{id} \;=\; [k,k^\perp] \circ [k,k^\perp]^\dagger \;&=\; [k,k^\perp] \circ \langle k^\dagger, (k^\perp)^\dagger\rangle \\ &=\; \nabla \circ \langle k \circ k^\dagger, k^\perp \circ (k^\perp)^\dagger\rangle \;=\; \nabla \circ p_k. \end{aligned}$$

4 Effect algebra structure on predicates

We start this section by introducing a partially defined sum $\ovee$ on predicates. This generalises the approach from [35], which is described only for maps of the form $X \to 1+1$. We show that such sums $\ovee$ turn the collection of (positive) predicates on an object into what will be described as an "effect algebra" in the next section. Such effect algebras are partially ordered.

Definition 3. *Let* ***C*** *be a category in which Assumption 1.(ii) holds: the two maps* $[\mathrm{id}, \kappa_2], [[\kappa_2, \kappa_1], \kappa_2]\colon (X+X)+X \rightrightarrows X+X$ *are jointly monic. We call two predicates* $p, q\colon X \to X+X$ *in* ***C*** orthogonal*, and write* $p \perp q$ *in that case, if there is a "bound" map* $b\colon X \to (X+X)+X$ *such that the following triangles commute.*

$$\begin{tikzcd} & X \arrow[dl, "p"'] \arrow[d, "b"] \arrow[dr, "q"] & \\ X+X & (X+X)+X \arrow[l, "{[\mathrm{id},\kappa_2]}"] \arrow[r, "{[[\kappa_2,\kappa_1],\kappa_2]}"'] & X+X \end{tikzcd}$$

Notice that there is at most one such bound, by Assumption 1.(ii).

Given a bound b for p, q we write $p \ovee q = (\nabla + \mathrm{id}) \circ b \colon X \to X + X$ for their sum. Thus, $\ovee$ is a partial, binary operation.

If p, q are positive predicates, we define $p \perp q$ and $p \ovee q$ as before, with the restriction that the bound b should be a 3-tuple of positive maps.

It is easy to see that $p \ovee q$ is again a predicate, since:

$$\begin{aligned} \nabla \circ (p \ovee q) \;=\; \nabla \circ (\nabla + \mathrm{id}) \circ b &= \nabla \circ [\kappa_1 \circ \nabla, \kappa_2] \circ b \\ &= [\nabla, \mathrm{id}] \circ b \\ &= \nabla \circ [\mathrm{id}, \kappa_2] \circ b \;=\; \nabla \circ p \;=\; \mathrm{id}. \end{aligned}$$

Example 7. We briefly investigate the predicates from Section 3.

(i). For two predicates $U, V \in \mathit{Pred}(X) \cong \mathcal{P}(X)$ in **Sets** the sum exists iff U, V are disjoint. In that case one takes as bound for the corresponding predicates $p_U, p_V \colon X \to X + X$ the map $b \colon X \to (X + X) + X$ given by:

$$b(x) \;=\; \begin{cases} \kappa_1\kappa_1 x & \text{if } x \in U \\ \kappa_1\kappa_2 x & \text{if } x \in V \\ \kappa_2 x & \text{if } x \notin U \text{ and } x \notin V. \end{cases}$$

This is well-defined since $U \cap V = \emptyset$. Then:

$$\begin{aligned} \big([\mathrm{id}, \kappa_2] \circ b\big)(x) \;&=\; \left\{ \begin{array}{ll} \kappa_1 x & \text{if } x \in U \\ \kappa_2 x & \text{if } x \notin U \end{array} \right\} \;=\; p_U(x) \\ \big([[\kappa_2, \kappa_1], \kappa_2] \circ b\big)(x) \;&=\; \left\{ \begin{array}{ll} \kappa_1 x & \text{if } x \in V \\ \kappa_2 x & \text{if } x \notin V \end{array} \right\} \;=\; p_V(x). \end{aligned}$$

The resulting join $p_U \ovee p_V$ is $p_{U \cup V}$ since:

$$\big((\nabla + \mathrm{id}) \circ b\big)(x) \;=\; \begin{cases} \kappa_1 x & \text{if } x \in U \text{ or } x \in V, \textit{i.e. } x \in U \cup V \\ \kappa_2 x & \text{if } x \notin U \text{ and } x \notin V, \textit{i.e. } x \notin U \cup V. \end{cases}$$

(ii). For the predicates $\mathit{Pred}(X) \subseteq \mathcal{P}(X) \times \mathcal{P}(X)$ on an object (set) X in the Kleisli category $\mathcal{K}\ell(\mathcal{P})$ of sets and relations the description of $\ovee$ above does not lead to a unique sum, since the "jointly monic" requirement does not hold in $\mathcal{K}\ell(\mathcal{P})$, as shown in Remark 2. Hence our predicate logic does not work for non-deterministic computation.

(iii). Next we consider two fuzzy predicates $\varphi, \psi \colon X \to [0, 1]$, with associated predicates $p_\varphi, p_\psi \colon X \to \mathcal{D}(X + X)$ in the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of sets and stochastic relations, as in Example 3. We claim that a bound exists iff $\varphi(x) + \psi(x) \leq 1$, for all $x \in X$. In that case one defines $b \colon X \to \mathcal{D}((X + X) + X)$ as:

$$b(x) \;=\; \varphi(x)\kappa_1\kappa_1 x + \psi(x)\kappa_1\kappa_2 x + (1 - \varphi(x) - \psi(x))\kappa_2 x.$$

The resulting sum $p_\varphi \ovee p_\psi$ corresponds to the pointwise sum $\varphi + \psi$, since:

$$\begin{aligned}\big(p_\varphi \ovee p_\psi\big)(x) &= \big((\nabla + \mathrm{id}) \circ b\big)(x) \\ &= (\varphi(x) + \psi(x))\kappa_1 x + (1 - \varphi(x) - \psi(x))\kappa_2 x.\end{aligned}$$

(iv). Let **C** be dagger biproduct category. We have already seen that in such a category a predicate $p = \langle p_1, p_2\rangle \colon X \to X \oplus X$ consists of a pair of maps $p_1, p_2 \colon X \rightrightarrows X$ with $p_1 + p_2 = \mathrm{id}$, where $+$ is the sum from (3).

If $b = \langle\langle b_1, b_2\rangle, b_3\rangle \colon X \to (X \oplus X) \oplus X$ is a bound for $p = \langle p_1, p_2\rangle \colon X \to X \oplus X$ and $q = \langle q_1, q_2\rangle \colon X \to X \oplus X$, then $b_1 = p_1$, $b_2 = q_1$ and so:

$$p \ovee q = \langle p_1 + q_1, b_3\rangle \qquad \text{where} \qquad p_1 + q_1 + b_3 = \mathrm{id}.$$

Hence this sum is a pair of positive maps.

In the category **BifMRel** this $\ovee$ involves the element-wise sum of positive predicates (as matrices), which is defined if the sum of the diagonal entries exists in $[0, 1]$, see Example 4.

In the category **Hilb** the first part p_1 of a positive predicate $p = \langle p_1, p_2\rangle$ is an effect. The previous observations show that the sum $\ovee$ on effects is the ordinary (pointwise) sum, if it is below the identity map:

$$\begin{aligned}\langle p_1, p_2\rangle \ovee \langle q_1, q_2\rangle &= \langle p_1 + q_1, \mathrm{id} - p_1 - q_1\rangle \qquad \text{assuming } p_1 + q_1 \leq \mathrm{id} \\ &= \langle p_1 + q_1, p_2 + q_2 - \mathrm{id}\rangle.\end{aligned}$$

We list some general results about the partial sum $\ovee$ on predicates. In the next section we see how these results fit in the general theory of effect algebras.

Lemma 4. *Let **C** be a category in which the "jointly monic" requirement from Assumption 1.(ii) holds. The resulting partially defined operation $\ovee$ on predicates:*

(i). is commutative: if $p \perp q$, then $q \perp p$ and $p \ovee q = q \ovee p$;

(ii). has $0 = \kappa_2$ has neutral element: $0 \perp p$ holds and $0 \ovee p = p$;

(iii). is associative if diagram (E) in Assumption 1.(i) is a pullback; this means that if $p \perp q$ and $p \ovee q \perp r$, then $q \perp r$ and $p \perp q \ovee r$, and moreover, $(p \ovee q) \ovee r = p \ovee (q \ovee r)$;

(iv). satisfies $p \ovee p^\perp = 1$;

(v). if diagram (K) is a pullback, then $p^\perp$ is the sole predicate q with $p \ovee q = 1$;

(vi). if diagram (K+) is a pullback, then $1 \perp p$ implies $p = 0$.

Proof. (i). The operation $\ovee$ is commutative, since if b is a bound for p, q, then $b' = ([\kappa_2, \kappa_1] + \mathrm{id}) \circ b$ is a bound for q, p:

$$\begin{aligned}[\mathrm{id}, \kappa_2] \circ b' &= [\mathrm{id}, \kappa_2] \circ ([\kappa_2, \kappa_1] + \mathrm{id}) \circ b = [[\kappa_2, \kappa_1], \kappa_2] \circ b = q \\ [[\kappa_2, \kappa_1], \kappa_2] \circ b' &= [[\kappa_2, \kappa_1], \kappa_2] \circ ([\kappa_2, \kappa_1] + \mathrm{id}) \circ b \\ &= [[\kappa_2, \kappa_1] \circ [\kappa_2, \kappa_1], \kappa_2] \circ b = [\mathrm{id}, \kappa_2] \circ b = p.\end{aligned}$$

This b and b' yield the same sum:

$$\begin{aligned} q \ovee p \;=\; (\nabla + \mathrm{id}) \circ b' \;&=\; (\nabla + \mathrm{id}) \circ ([\kappa_2, \kappa_1] + \mathrm{id}) \circ b \\ &=\; ((\nabla \circ [\kappa_2, \kappa_1]) + \mathrm{id}) \circ b \;=\; (\nabla + \mathrm{id}) \circ b \;=\; p \ovee q. \end{aligned}$$

(ii). The zero predicate $0 = \kappa_2 \colon X \to X + X$ is neutral element for $\ovee$: the equation $0 \ovee p = p$ holds via the bound $b = (\kappa_2 + \mathrm{id}) \circ p \colon X \to (X + X) + X$.

(iii). For associativity assume predicates $p, q, r \colon X \to X + X$ where $p \perp q$, say via bound a, and $(p \ovee q) \perp r$, via bound b. Thus we have equations:

$$\left\{ \begin{aligned} [\mathrm{id}, \kappa_2] \circ a \;&=\; p \\ [[\kappa_2, \kappa_1], \kappa_2] \circ a \;&=\; q \\ (\nabla + \mathrm{id}) \circ a \;&=\; p \ovee q \end{aligned} \right. \qquad \left\{ \begin{aligned} [\mathrm{id}, \kappa_2] \circ b \;&=\; p \ovee q \\ [[\kappa_2, \kappa_1], \kappa_2] \circ b \;&=\; r \\ (\nabla + \mathrm{id}) \circ b \;&=\; (p \ovee q) \ovee r. \end{aligned} \right.$$

We consider the following pullback, which is an instance of (E) in Assumption 1,

$$\begin{tikzcd}
X \arrow[d, "b"'] \arrow[dr, dashed, "c"] \arrow[drr, bend left, "a"] & & \\
(X+X)+X \arrow[dr, "\alpha^{-1}"', "\cong"] & (X+X)+(X+X) \arrow[r, "\mathrm{id}+\nabla"] \arrow[d, "\nabla+\mathrm{id}"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "(E)"] & (X+X)+X \arrow[d, "\nabla+\mathrm{id}"] \\
& X+(X+X) \arrow[r, "\mathrm{id}+\nabla"'] & X+X
\end{tikzcd}$$

where $\alpha = [\kappa_1 \circ \kappa_1, \kappa_2 + \mathrm{id}]$ is the obvious associativity isomorphism.

We now take $c' = [[\kappa_2, \kappa_1 \circ \kappa_1], \kappa_2 + \mathrm{id}] \circ c \colon X \to (X + X) + X$. This c' is a bound for q and r, giving $q \perp r$. Next, the map $c'' = [\kappa_1, \kappa_2 + \mathrm{id}] \circ c \colon X \to (X + X) + X$ is a bound for p and $q \ovee r = (\nabla + \mathrm{id}) \circ c'$. Finally we obtain the associativity of $\ovee$:

$$\begin{aligned} p \ovee (q \ovee r) \;=\; (\nabla + \mathrm{id}) \circ c'' \;&=\; (\nabla + \mathrm{id}) \circ [\kappa_1, \kappa_2 + \mathrm{id}] \circ c \\ &=\; [[\kappa_1, \kappa_1], [\kappa_1, \kappa_2]] \circ c \\ &=\; [[\kappa_1, \kappa_1], \kappa_2] \circ [\kappa_1 \circ \kappa_1 \circ \nabla, \kappa_2 + \mathrm{id}] \circ c \\ &=\; [[\kappa_1, \kappa_1], \kappa_2] \circ \alpha \circ (\nabla + \mathrm{id}) \circ c \\ &=\; (\nabla + \mathrm{id}) \circ b \;=\; (p \ovee q) \ovee r. \end{aligned}$$

(iv). It is not hard to see that $b = \kappa_1 \circ p \colon X \to (X + X) + X$ is a bound for p and $p^\perp = p \circ [\kappa_2, \kappa_1]$, yielding $p \ovee p^\perp = 1$.

(v). Next we show that $p^\perp$ is the only predicate with $p \ovee p^\perp = 1$. So assume also for $q \colon X \to X + X$ we have $p \ovee q = 1$, say via bound b. Then $[\mathrm{id}, \kappa_2] \circ b = p$, $[[\kappa_2, \kappa_1], \kappa_2] \circ b = q$ and $(\nabla + \mathrm{id}) \circ b = 1 = \kappa_1$. We use a version of pullback (K) from Assumption 1 in:

$$\begin{tikzcd}
X \arrow[dr, dashed, "c"] \arrow[drr, bend left, "b"] \arrow[ddr, equal, bend right] & & \\
& X+X \arrow[r, "\kappa_1"] \arrow[d, "\nabla"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "(K)"] & (X+X)+X \arrow[d, "\nabla+\mathrm{id}"] \\
& X \arrow[r, "\kappa_1"'] & X+X
\end{tikzcd}$$

The fact that the bound b is of the form $\kappa_1 \circ c$ is enough to obtain $q = p^{\perp}$:

$$\begin{aligned} p^{\perp} \;=\; [\kappa_2, \kappa_1] \circ p \;=\; [\kappa_2, \kappa_1] \circ [\mathrm{id}, \kappa_2] \circ b &= [[\kappa_2, \kappa_1], \kappa_1] \circ \kappa_1 \circ c \\ &= [[\kappa_2, \kappa_1], \kappa_2] \circ \kappa_1 \circ c \\ &= [[\kappa_2, \kappa_1], \kappa_2] \circ b \;=\; q. \end{aligned}$$

(vi). Finally, suppose $1 \perp p$; we must prove $p = 0$. We may assume a bound b with $[\mathrm{id}, \kappa_2] \circ b = 1 = \kappa_1$ and $[[\kappa_2, \kappa_1], \kappa_2] \circ b = p$. Now we use the diagram (K+) to obtain a mediating map $X \to X$, that must be the identity, in:

$$\begin{tikzcd}
X \arrow[rr, "b"] \arrow[dr, dashed, "\mathrm{id}"] \arrow[ddr, equal, bend right] & & (X+X)+X \arrow[dr, "\alpha^{-1}", "\cong"'] \arrow[ddr, bend left=60, "{[\mathrm{id},\kappa_2]}"] & \\
& X \arrow[rr, "\kappa_1"] \arrow[d, equal] \arrow[d, phantom, "\lrcorner", very near start, shift left=2] \arrow[drr, phantom, "(K+)"] & & X+(X+X) \arrow[d, "\mathrm{id}+\nabla"] \\
& X \arrow[rr, "\kappa_1"'] & & X+X
\end{tikzcd}$$

As a result, $b = \alpha \circ \kappa_1 = \kappa_1 \circ \kappa_1$. Then:

$$p \;=\; [[\kappa_2, \kappa_1], \kappa_2] \circ b \;=\; [[\kappa_2, \kappa_1], \kappa_2] \circ \kappa_1 \circ \kappa_1 \;=\; \kappa_2 \;=\; 0. \qquad \square$$

5 Intermezzo on effect algebras

In the previous section we have seen that on predicates $X \to X+X$ a partial sum $\ovee$ can be defined — assuming the coproduct $+$ satisfies some basic properties. This structure will be described more abstractly in terms of effect algebras. This section provides a crash course for the relevant theory.

In order to define an effect algebra, we need the notion of partial commutative monoid (PCM). Before reading the definition of PCM, think of the unit interval $[0, 1]$ with addition $+$. This $+$ is obviously only a partial operation, which is commutative and associative in a suitable sense. This will be formalised next.

Definition 4. *A partial commutative monoid (PCM) consists of a set M with a zero element $0 \in M$ and a partial binary operation $\ovee\colon M \times M \to M$ satisfying the three requirements below. They involve the notation $x \perp y$ for: $x \ovee y$ is defined; in that case x, y are called orthogonal.*

(i). Commutativity: $x \perp y$ implies $y \perp x$ and $x \ovee y = y \ovee x$;
(ii). Associativity: $y \perp z$ and $x \perp (y \ovee z)$ implies $x \perp y$ and $(x \ovee y) \perp z$ and also $x \ovee (y \ovee z) = (x \ovee y) \ovee z$;
(iii). Zero: $0 \perp x$ and $0 \ovee x = x$;

The first three points of Lemma 4 say that predicates $p\colon X \to X+X$ form a PCM, if the jointly monic requirement from Assumption 1 holds. But also, for instance, for each set X the lift $\{0\} + X$ of X, obtained by adjoining a new element 0, is an example of a PCM, with $u \ovee 0 = u = 0 \ovee u$, and $\ovee$ undefined otherwise. Such structures are also studied under the name ‘partially additive monoid’ in program semantics, see [3].

The notion of effect algebra extends a PCM with an orthocomplement $(-)^{\perp}$. It is due to [21]; the same notion has been introduced in [13] and in [23]. The reference [20] gives an overview.

Definition 5. *An effect algebra is a PCM $(E, 0, \ovee)$ with an orthocomplement. The latter is a unary operation $(-)^{\perp} \colon E \to E$ satisfying:*

- *$x^{\perp} \in E$ is the unique element in E with $x \ovee x^{\perp} = 1$, where $1 = 0^{\perp}$;*
- *$x \perp 1 \Rightarrow x = 0$.*

For such an effect algebra one defines,

$$x \owedge y = (x^{\perp} \ovee y^{\perp})^{\perp} \qquad \textit{and} \qquad \begin{cases} x \leq y \iff \exists_z .\, x \ovee z = y \\ y \ominus x = z \iff y = x \ovee z. \end{cases}$$

We shall shortly see, in point (iii) of Lemma 5, that there is at most one element z with $x \ovee z = y$.

*A homomorphism $E \to D$ of effect algebras is given by a function $f \colon E \to D$ between the underlying sets satisfying $f(1) = 1$, and if $x \perp x'$ in E then both $f(x) \perp f(x')$ in D and $f(x \ovee x') = f(x) \ovee f(x')$. Effect algebras and their homomorphisms form a category, written as **EA**.*

When writing $x \ovee y$ we shall implicitly assume that $x \ovee y$ is defined, *i.e.* $x \perp y$. Similarly for $\owedge$ and $\ominus$. The notation $\ovee$ might suggest that this operation is idempotent, but this is not the case (and not intended): for instance, in $[0, 1]$ one as $x \ovee x \neq x$ if $0 < x \leq \frac{1}{2}$.

Example 8. We briefly discuss several (classes of) examples of effect algebras.

(i). A singleton set forms an example of a degenerate effect algebra, with $0 = 1$; it is final in the category **EA**. A two element set $2 = \{0, 1\}$ is initial in **EA**.

(ii). A more interesting example is the unit interval $[0, 1] \subseteq \mathbb{R}$ of real numbers, with $r^{\perp} = 1 - r$ and $r \ovee s$ is defined as $r + s$ in case this sum is in $[0, 1]$. In fact, for each non-negative number $M \in \mathbb{R}$ the interval $[0, M] = \{r \in \mathbb{R} \mid 0 \leq r \leq M\}$ is an example of an effect algebra, with $r^{\perp} = M - r$.

Also the interval $[0, M] \cap \mathbb{Q}$ of rational numbers, for non-negative $M \in \mathbb{Q}$, is an effect algebra. And so is the interval $[0, M] \cap \mathbb{N}$ of natural numbers, for $M \in \mathbb{N}$.

The general situation involves so-called “interval effect algebras”, see *e.g.* [22] or [20, 1.4]. An Abelian group $(G, 0, -, +)$ is called ordered if it carries a partial order $\leq$ such that $a \leq b$ implies $a + c \leq b + c$, for all $a, b, c \in G$. A positive point is an element $p \in G$ with $p \geq 0$. For such a point the interval $[0, p] = \{a \in G \mid 0 \leq a \leq p\}$ forms an effect algebra with p as top, orthocomplement $a^{\perp} = p - a$, and sum $a + b$, which is considered to be defined in case $a + b \leq p$.

(iii). A separate class of examples has a join as sum $\ovee$. Let $(L, \vee, 0, (-)^{\perp})$ be an ortholattice: $\vee, 0$ are finite joins and complementation $(-)^{\perp}$ satisfies $x \leq y \Rightarrow y^{\perp} \leq x^{\perp}$, $x^{\perp\perp} = x$ and $x \vee x^{\perp} = 1 = 0^{\perp}$. This L is called an orthomodular lattice if $x \leq y$

implies $y = x \vee (x^{\perp} \wedge y)$. Such an orthomodular lattice forms an effect algebra in which $x \ovee y$ is defined if and only if $x \perp y$ (*i.e.* $x \leq y^{\perp}$, or equivalently, $y \leq x^{\perp}$); and in that case $x \ovee y = x \vee y$. This restriction of $\vee$ is needed for the validity of requirements (1) and (2) in Definition 5:

- suppose $x \ovee y = 1$, where $x \perp y$, *i.e.* $x \leq y^{\perp}$. Then, by the orthomodularity property, we get uniqueness of orthocomplements:

$$y^{\perp} = x \vee (x^{\perp} \wedge y^{\perp}) = x \vee (x \vee y)^{\perp} = x \vee 1^{\perp} = x \vee 0 = x.$$

- $x \perp 1$ means $x \leq 1^{\perp} = 0$, so that $x = 0$.

In particular, the lattice $\mathit{KSub}(H)$ of closed linear subspaces of a Hilbert space H is an orthomodular lattice and thus an effect algebra. This applies more generally to the kernel subobjects of an object in a dagger kernel category [30].

(iv). Since Boolean algebras are (distributive) orthomodular lattices, they are also effect algebras. They satisfy $x \perp y$ iff $x \wedge y = 0$, see the proof of Lemma 8 below.

(v). For an arbitrary set A write $\mathit{MO}(A) = (1 + A) + (1 + A) = (A + A) \cup \{0, 1\}$, which can be understood as a poset with top and bottom and two copies of A in between so that one can define an orthocomplement $(-)^{\perp}$ on $\mathit{MO}(A)$, by swapping. On this $\mathit{MO}(A)$ one can actually define $\ovee$ in a minimal way, namely as:

$$z \ovee w = \begin{cases} z \;\; \text{if } w = 0 \\ w \;\; \text{if } z = 0 \\ 0 \;\; \text{if } z = 1 \text{ or } w = 1 \\ 1 \;\; \text{if } z, w \text{ are equal elements in different } A\text{-components} \\ \text{undefined, otherwise.} \end{cases}$$

There is an obvious insertion function $A \to \mathit{MO}(A)$, that makes $\mathit{MO}(A) \in \mathbf{EA}$ the free effect algebra on the set A.

We summarise some properties of effect algebras.

Lemma 5. *In an effect algebra one has:*

(i). *Orthocomplementing is an involution:* $x^{\perp\perp} = x$;
(ii). *Top and bottom are each others orthocomplements:* $1^{\perp} = 0$ *and* $0^{\perp} = 1$;
(iii). *Cancellation holds:* $x \ovee y = x \ovee z$ *implies* $y = z$;
(iv). *Positivity (or zerosumfreeness) holds:* $x \ovee y = 0$ *implies* $x = y = 0$;
(v). $\leq$ *is a partial order with* 1 *as top and* 0 *as bottom element;*
(vi). $x \leq y$ *implies* $x \ovee z \leq y \ovee z$;
(vii). $x \leq y$ *implies* $y^{\perp} \leq x^{\perp}$;
(viii). $x \ovee y$ *is defined iff* $x \perp y$ *iff* $y \leq x^{\perp}$ *iff* $x \leq y^{\perp}$;
(ix). $\owedge$ *with 1 also forms a partial commutative monoid structure.*

Proof. For point (i) we use $x \ovee x^{\perp} = 1 = x^{\perp\perp} \ovee x^{\perp}$, so that $x = x^{\perp\perp}$ since the orthocomplement of $x^{\perp}$ is unique. Uniqueness also yields $1^{\perp} = 0$ from $1 \ovee 1^{\perp} = 1 = 1 \ovee 0$. Hence $0^{\perp} = 1^{\perp\perp} = 1$.

For cancellation, assume $x \ovee y = x \ovee z$. The orthocomplement $u = (x \ovee y)^{\perp} = (x \ovee z)^{\perp}$ satisfies $x \ovee y \ovee u = 1 = x \ovee z \ovee u$, so that $y = (x \ovee u)^{\perp} = z$ by uniqueness of orthocomplements. If $x \ovee y = 0$ in (iv), then $y^{\perp} = 0 \ovee y^{\perp} = x \ovee y \ovee y^{\perp} = x \ovee 1$. Hence $x \perp 1$ and thus $x = 0$. Similarly $y = 0$.

For (v) we obtain reflexivity $x \leq x$ from $x \ovee 0 = x$. Transitivity is easy: if $x \leq y$ and $y \leq z$, say via $y = x \ovee u$ and $z = y \ovee v$, then $z = y \ovee v = x \ovee u \ovee v$, so that $x \leq z$. Finally, if $x \leq y$ and $y \leq x$, say via $y = x \ovee u$ and $x = y \ovee v$, then $x \ovee 0 = x = x \ovee u \ovee v$. By cancelling x we get $0 = u \ovee v$, so that $u = v = 0$ by positivity, and thus $x = y$. We get $0 \leq x$ since $0 \ovee x = x$, and $x \leq 1$ since $x \ovee x^{\perp} = 1$.

For (vi) and (vii) assume $x \leq y$, say via $y = x \ovee u$. Then $x \ovee z \ovee u = x \ovee u \ovee z = y \ovee z$, so that $x \ovee z \leq y \ovee z$. Also, $1 = y \ovee y^{\perp} = x \ovee u \ovee y^{\perp}$, so that $x^{\perp} = u \ovee y^{\perp}$, showing that $y^{\perp} \leq x^{\perp}$.

For (viii) first assume $x \perp y$. Then $x \ovee y \ovee (x \ovee y)^{\perp} = 1$, so that $x^{\perp} = y \ovee (x \ovee y)^{\perp}$, and thus $y \leq x^{\perp}$. Conversely, if $x^{\perp} = y \ovee u$, then $x \ovee y \ovee u = 1$, so that $x \ovee y$ exists.

In (ix) it is easy to see that $\owedge$ is commutative and associative, as partial operation. It has 1 as unit, since $x \owedge 1 = (x^{\perp} \ovee 1^{\perp})^{\perp} = (x^{\perp} \ovee 0)^{\perp} = x^{\perp\perp} = x$. $\square$

The minus (or difference) operation $\ominus$ introduced in Definition 5 satisfies many expected properties. We only list them here and leave the proof to the interested reader.

Lemma 6. *In an effect algebra:*

(i). $x \ominus y$ *is defined if and only if* $y \leq x$*;*
(ii). $x \ominus 0 = x$ *and* $1 \ominus x = x^{\perp}$*;*
(iii). $x \ominus y = (x^{\perp} \ovee y)^{\perp} = x \owedge y^{\perp}$*;*
(iv). $x \ovee y \leq z$ *iff* $x \leq z \ominus y$*;*
(v). $z \ominus x = z \ominus y$ *implies* $x = y$*;*
(vi). $(x \ovee y) \ominus y = x$*;*
(vii). $x \leq y$ *implies* $y \ominus x \leq y$ *and* $y \ominus (y \ominus x) = x$*;*
(viii). $x \leq y \leq z$ *implies* $z \ominus y \leq z \ominus x$ *and* $(z \ominus x) \ominus (z \ominus y) = y \ominus x$*;*
(ix). $x \ominus y \leq z$ *iff* $x \leq z \ovee y$*.* $\square$

From points (iv) and (ix) we may conclude that $- \ominus y$, considered as functor $\uparrow y \rightarrow \downarrow y^{\perp}$ between posets, has $- \ovee y \colon \downarrow y^{\perp} \rightarrow \uparrow y$ both as left and right adjoint: $- \ovee y \dashv - \ominus y \dashv - \ovee y$. Hence $- \ominus y$ and $- \ovee y$ preserve all meets and joins that exist in $\uparrow y$ and $\downarrow y^{\perp}$, respectively.

Notice that in the context of orthomodular lattices, for $y \leq x$ one has $x \ominus y = x \wedge y^{\perp}$. Since $x \ominus y \leq y^{\perp}$ one has $x \ominus y \perp y$, so that $(x \ominus y) \ovee y = (x \wedge y^{\perp}) \vee y = x$, the latter by orthomodularity.

We can now draw appropriate conclusions about the structure on predicates as investigated in the previous section.

Proposition 1. *Let* ***C*** *be a category with coproducts* $+$ *and let* $X \in$ ***C*** *be an arbitrary object.*

*(i). If these coproducts satisfy requirements (i) and (ii) from Assumption 1, then the predicates Pred(X) on $X \in$ **C** form an effect algebra.*
*(ii). If **C** is a dagger additive category which is zerosumfree, then the positive predicates pPred(X) on X also form an effect algebra.*

Proof. The first point follows directly from Lemma 4. For the second point we have to check carefully that Lemma 2 applies in the situation where we use the pullback (K+) in the proof of Lemma 4: the relevant map g in Assumption 1.(i) must be zero monic. In the final diagram in the proof of Lemma 4 this map g is the codiagonal $\nabla\colon X \oplus X \to X$. The zerosumfreeness requirement implies that ∇ is zero monic:

$$\nabla \circ \langle b_2, b_3 \rangle = b_2 + b_3 = 0 \Longrightarrow b_2 = b_3 = 0 \Longrightarrow \langle b_2, b_3 \rangle = 0,$$

where we use that the maps b_i from bound b are positive (by definition). □

The first point applies directly to each extensive category, by Lemma 1, and to the Kleisli category $\mathcal{K}\ell(\mathcal{D})$, by Lemma 3. The second point applies for instance to the category **Hilb** of Hilbert spaces. In all of these cases the induced ordering on predicates is the expected one. For instances, it is the pointwise order on fuzzy predicates in $[0,1]^X$. On the effects $\mathcal{E}\!f(H)$ of a Hilbert space H we get the so-called Löwner partial order: $A \leq B$ iff there is a positive operator $C\colon H \to H$ with $A + C = B$.

5.1 Effect algebra homomorphisms

Homomorphisms of effect algebras (see Definition 5) are like finitely additive probability measures. Homomorphisms $E \to [0,1]$ to the unit interval are sometimes called states. They form a convex subset, see Subsection 5.3 below. Effect algebra maps preserve all the relevant structure.

Lemma 7. *Let $f\colon E \to D$ be a homomorphism of effect algebras. Then:*

$$f(x^{\perp}) = f(x)^{\perp} \quad \textit{and thus} \quad f(0) = 0,$$

and thus also, when the left-hand-sides inside $f(-)$ are defined:

$$f(x \owedge x') = f(x) \owedge f(x') \quad \textit{and} \quad f(x \ominus x') = f(x) \ominus f(x'),$$

and:

$$x \leq x' \Longrightarrow f(x) \leq f(x').$$

Proof. From $1 = f(1) = f(x \ovee x^{\perp}) = f(x) \ovee f(x^{\perp})$ we obtain $f(x^{\perp}) = f(x)^{\perp}$ by uniqueness of orthocomplements. In particular, $f(0) = f(1^{\perp}) = f(1)^{\perp} = 1^{\perp} = 0$. □

Example 9. Orthocomplement $(-)^{\perp}$ is a homomorphism $E \to E^{\mathrm{op}}$ in **EA**, namely from $(E, 0, \ovee, (-)^{\perp})$ to $(E, 1, \owedge, (-)^{\perp})$.

Using the free construction *MO* from Example 8.(v), an element (or point) $x \in E$ of an effect algebra E can be identified with a homomorphism *MO*$(1) = 2 \times 2 \to E$ in **EA**, as in:



On the homset **EA**(E, D) of maps $E \to D$ in **EA** one may define $(-)^\perp$ and $\ovee$ pointwise, as in $f^\perp(x) = f(x)^\perp$. But this does not yield a homomorphism $E \to D$, since for instance $f^\perp(1) = f(1)^\perp = 1^\perp = 0$. Hence these homsets do not form effect algebras.

The Kochen-Specker Theorem [42] can be formulated in terms of homomorphisms of effect algebras, like in [17, §§5.1], namely as: there is no effect algebra morphism *KSub*$(H) \to 2 = \{0, 1\}$, from the closed linear subspaces of a Hilbert space H, with $\dim(H) \geq 3$, to the two-element (Boolean / initial) effect algebra.

Recall that homomorphisms of effect algebras preserve only sums of orthogonal elements. For Boolean algebras it turns out that this implies preservation of all sums (or joins).

Lemma 8. *Let B, C be Boolean algebras (and thus effect algebras). Then: $f\colon B \to C$ is a homomorphism of effect algebras if and only if it is a homomorphism of Boolean algebras. As a result, the inclusion functor **BA** $\to$ **EA** from Boolean algebras to effect algebras is full and faithful.*

Proof. The first observation is that in a Boolean algebra $x \perp y$ iff $x \wedge y = 0$, since:

- if $x \perp y$, *i.e.* $x \leq \neg y$, one gets $x \wedge y \leq \neg y \wedge y = 0$.
- if $x \wedge y = 0$, then by distributivity: $x = x \wedge 1 = x \wedge \neg(x \wedge y) = x \wedge (\neg x \vee \neg y) = (x \wedge \neg x) \vee (x \wedge \neg y) = 0 \vee (x \wedge \neg y) = x \wedge \neg y$. Hence $x \leq \neg y$.

Next notice that the join of two elements x, y can be written disjointly as:

$$x \vee y \,=\, (x \wedge \neg y) \vee (x \wedge y) \vee (y \wedge \neg x). \tag{6}$$

If $f\colon B \to C$ is a morphism of effect algebras, it preserves disjoint joins, and by monotonicity it satisfies $f(x) \vee f(y) \leq f(x \vee y)$ and $f(x \wedge y) \leq f(x) \wedge f(y)$. Therefore, by (6),

$$\begin{aligned} f(x \vee y) \,&=\, f\big((x \wedge \neg y) \vee (x \wedge y) \vee (y \wedge \neg x)\big) \\ &=\, f(x \wedge \neg y) \vee f(x \wedge y) \vee f(y \wedge \neg x) \\ &\leq\, \big(f(x) \wedge \neg f(y)\big) \vee \big(f(x) \wedge f(y)\big) \vee \big(f(y) \wedge \neg f(x)\big) \\ &=\, f(x) \vee f(y). \end{aligned}$$

Hence f preserves joins — and also meets, since $x \wedge y = \neg(\neg x \vee \neg y)$ — and is thus morphism of Boolean algebras. □

We consider some obvious constructions on effect algebras.

Lemma 9. *Let E, D be effect algebras.*

*(i). The opposite $E^{op} = (E, 1, \owedge, (-)^\perp)$, with order $x \leq^{op} y$ iff $y \leq x$ is an effect algebra; this yields an involutive functor $(-)^{op}\colon$ **EA**$^{op} \to$ **EA**.*
*(ii). The product $E \times D$ with component-wise operations and obvious projections is a cartesian product in **EA**.*

(iii). The coproduct in **EA** *is given by* $E + D = \big((E - \{0,1\}) + (D - \{0,1\})\big) + \{0,1\}$*, where* $+$ *on the right-hand-side of the equality sign is the disjoint union (or coproduct) of sets, with obvious coprojections* $E \rightarrow E + D \leftarrow D$*.*

(iv). The downset $\downarrow x = \{y \in E \mid y \leq x\}$ *is an effect algebra, for* $x \in E$*.*

The coproduct effect algebra $E + D$ is obtained by taking the disjoint union of E, D but with their top and bottom elements $1, 0$ identified. This is sometimes called an amalgamated sum.

Proof. It is obvious that the opposite and product are effect algebras. For the coproduct we define for $z, w \in E + D$,

$$z^{\perp} = \begin{cases} 1 & \text{if } z = 0 \\ 0 & \text{if } z = 1 \\ x^{\perp} & \text{if } z = x \in E \\ y^{\perp} & \text{if } z = y \in D. \end{cases} \qquad z \ovee w = \begin{cases} w & \text{if } z = 0 \\ z & \text{if } w = 0 \\ x \ovee u & \text{if } z = x \in E, w = u \in E \\ y \ovee v & \text{if } z = y \in D, w = v \in D \\ \text{undefined, otherwise.} \end{cases}$$

A downset $\downarrow x$ is again an effect algebra with x as top, complement $y^{\perp_x} = x \ominus y$ and sum $y \ovee z$ defined iff $y \ovee z \leq x$. Then $y \ovee y^{\perp_x} = y \ovee (x \ominus y) = x$. □

5.2 Effect monoids

Abstractly, an effect monoid is a monoid in the category of effect algebras, just like a semiring is a monoid in the category of commutative monoids. An effect monoid is thus an effect algebra with a multiplication operation. In a Boolean algebra this is the meet $\wedge$, but for fuzzy predicates in $[0,1]^X$ it is pointwise multiplication.

In [36] it is shown that the category **EA** of effect algebras is symmetric monoidal, where morphisms $E \otimes D \rightarrow C$ in **EA** correspond to ‘bimorphisms’ $f \colon E \times D \rightarrow C$, satisfying $f(1,1) = 1$, and for all $x, x' \in E$ and $y, y' \in D$,

$$\begin{cases} x \perp x' \implies f(x,y) \perp f(x',y) \;\text{ and }\; f(x \ovee x', y) \;=\; f(x,y) \ovee f(x',y) \\ y \perp y' \implies f(x,y) \perp f(x,y') \;\text{ and }\; f(x, y \ovee y') \;=\; f(x,y) \ovee f(x,y'). \end{cases}$$

The tensor unit is the two-element (initial) effect algebra $2 = \{0,1\}$.

Example 10. For an arbitrary set X the powerset $\mathcal{P}(X)$ is a Boolean algebra, and so an orthomodular lattice, and thus an effect algebra. The tensor product $[0,1] \otimes \mathcal{P}(X)$ of effect algebras is given by the set of step functions $f \colon X \rightarrow [0,1]$; such functions have only finitely many output values. When X is a finite set, say with n elements, then $[0,1] \otimes \mathcal{P}(X) \cong [0,1]^n$, see [25].

As special case we have $[0,1] \otimes \{0,1\} \cong [0,1]$, since $\{0,1\}$ is the tensor unit. Writing *MO*(n) for the free effect algebra with $2n+2$ elements, like in Example 8.(v), we may write this as $[0,1] \otimes$ *MO*$(0) \cong [0,1]$. Next, $[0,1] \otimes$ *MO*$(1) \cong [0,1] \otimes \mathcal{P}(\{0,1\}) \cong [0,1]^2$. It can be shown that $[0,1] \otimes$ *MO*(2) is an octahedron.

Gleason's Theorem [24] gives an isomorphism:

$$\mathbf{EA}\big(\mathit{KSub}(H), [0,1]\big) \;\cong\; \mathcal{DM}(H),$$

between this homset of effect algebra maps, from the closed subsets of a Hilbert space H to the unit interval $[0,1]$, and the density matrices on H — assuming that H is of dimension at least 3. In [37] it is shown that Gleason's Theorem amounts to an isomorphism of effect algebras $[0,1] \otimes \mathit{KSub}(H) \;\cong\; \mathcal{E}f(H)$, relating projections and effects on a Hilbert space H.

Using this symmetric monoidal structure $(\otimes, 2)$ on **EA** we can consider, in a standard way, the categories **Mon**(**EA**) and **CMon**(**EA**) of (commutative) monoids in the category **EA** of effect algebras. A monoid $M \in$ **Mon**(**EA**) consists of a set M carrying effect algebra structure $(0, \ovee, (-)^{\perp})$ and a monoid structure, written multiplicatively, as in: $M \otimes M \overset{\cdot}{\to} M \leftarrow 2$. Since 2 is initial, the latter map $M \leftarrow 2$ does not add any structure. The monoid structure on M is thus determined by a bimorphism $\cdot\colon M \times M \to M$ that preserves $\ovee$ in each variable separately and satisfies $1 \cdot x = x = x \cdot 1$. The main example of a (commutative) effect monoid is the unit interval $[0,1]$, with ordinary multiplication. For $r_1 + r_2 \leq 1$ we have the familiar distributivity in each variable:

$$s \cdot (r_1 \ovee r_2) \;=\; s \cdot (r_1 + r_2) \;=\; (s \cdot r_1) + (s \cdot r_2) \;=\; (s \cdot r_1) \ovee (s \cdot r_2).$$

More generally, each set of fuzzy predicates $[0,1]^X$ is a commutative effect monoid, via pointwise multiplication. But also, each Boolean algebra forms a commutative effect monoid, with meet $\wedge$ as multiplication.

In Subsection 2.3 we have seen the distribution monad $\mathcal{D}(X)$ consisting of formal sums $\sum_i r_i x_i$ with $r_i \in [0,1]$ satisfying $\sum_i r_i = 1$. In [35] it is shown that such a distribution monad can also be defined wrt. a (commutative) effect monoid M instead of $[0,1]$. This works as follows.

$$\mathcal{D}_M(X) \;=\; \{m_1 x_1 + \cdots + m_k x_k \mid x_i \in X, m_i \in M \text{ with } \ovee_i\, m_i = 1 \in M\}.$$

Implicitly in this formulation we assume that the finite sum $\ovee_i\, m_i$ exists. It can be shown that $\mathcal{D}_M$ is a monad on **Sets**, just like $\mathcal{D} = \mathcal{D}_{[0,1]}$ is a monad.

Here we are interested in the category $\mathbf{Conv}_M = \mathcal{EM}(\mathcal{D}_M)$ of Eilenberg-Moore algebras of this monad $\mathcal{D}_M$. Its objects are "convex sets over M", that is, sets X in which for each formal convex sum $\sum_i m_i x_i$ an element $\ovee_i\, m_i x_i \in X$ exists. Morphisms in $\mathcal{EM}(\mathcal{D}_M)$ are *affine* maps; they preserves such convex sums $\ovee_i\, m_i x_i$.

Example 11. For $M = [0,1]$ the category $\mathbf{Conv}_{[0,1]}$ is the usual category of convex sets, see *e.g.* [56,57,33]. If we take the initial effect algebra as effect monoid $M = 2 = \{0,1\} = \mathcal{P}(1)$ we see that $\mathcal{D}_2(X) \cong X$, and that the associated category of convex sets $\mathbf{Conv}_2$ is simply the category **Sets** of ordinary sets.

5.3 Effect modules

For an effect monoid $M \in$ **Mon**(**EA**) we can consider the category $\mathbf{Act}_M(\mathbf{EA}) = \mathbf{EMod}_M$ of M-monoid actions (scalar multiplications), or *effect modules* over M (see

[48, VII,§4]). This resembles the category $\mathbf{Mod}_S$ of modules over a semiring S which may be described as the category $\mathbf{Act}_S(\mathbf{CMon})$ of commutative monoids with S-scalar multiplication. An effect module $E \in \mathbf{Act}_M(\mathbf{EA})$ thus consists of an effect algebra E together with an action (or scalar multiplication) $\bullet\colon M \otimes E \to E$, corresponding to a bimorphism $M \times E \to E$. A homomorphism of effect modules $E \to D$ consists of a map of effect algebras $f\colon E \to D$ preserving scalar multiplication $f(m \bullet x) = m \bullet f(x)$ for all $m \in M$ and $x \in E$.

Example 12. The main example of a (commutative) monoid in **EA** is the unit interval $[0,1] \in \mathbf{EA}$ via ordinary multiplication. The associated effect modules $\mathbf{EMod}_{[0,1]} = \mathbf{Act}_{[0,1]}(\mathbf{EA})$ have been studied earlier under the name 'convex effect algebras', see [53]. Convex sums exist since if $r \perp s$ in an effect monoid M, then the sum $rx \ovee sy$ exists for all elements x, y in an effect module over M.

The effects $\mathcal{E}f(H)$ of a Hilbert space form an example of an effect module over $[0,1]$, with the usual scalar multiplication $[0,1] \times \mathcal{E}f(H) \to \mathcal{E}f(H)$.

We can also consider effect modules over the effect monoid $2 = \{0,1\}$. Since $2 \in \mathbf{EA}$ is the tensor unit, the scalar multiplication becomes trivial. Hence we have an isomorphism $\mathbf{EMod}_2 \cong \mathbf{EA}$.

In [38] a (dual) adjunction between convex sets over $[0,1]$ and effect modules over $[0,1]$ is described. This adjunction exists in fact for an arbitrary effect monoid M — instead of $[0,1]$ — by using M as dualising object. It gives translations back-and-forth between Heisenberg's and Schrödinger's view on quantum computation.

Proposition 2. *Let M be an effect monoid. By "homming into M" one obtains an adjunction:*

$$\begin{tikzcd}
\big(\boldsymbol{EMod}_M\big)^{op} \arrow[r, bend left, "\boldsymbol{Conv}(-{,}M)"] \arrow[r, phantom, "\top"] & \boldsymbol{Conv}_M = \mathcal{EM}(\mathcal{D}_M) \arrow[l, bend left, "\boldsymbol{EMod}(-{,}M)"]
\end{tikzcd}$$

Proof. Given a convex set $X \in \mathbf{Conv}_M$, the homset $\mathbf{Conv}(X, M)$ of affine maps is an effect module, with $f \perp g$ iff $\forall_{x \in X}.\, f(x) \perp g(x)$ in M. In that case one defines $(f \ovee g)(x) = f(x) \ovee g(x)$. It is easy to see that this is again an affine function. Similarly, the pointwise scalar product $(m \bullet f)(x) = m \cdot f(x)$ yields an affine function. This mapping $X \mapsto \mathbf{Conv}(X, M)$ gives a contravariant functor since for $h\colon X \to X'$ in $\mathbf{Conv}_M$ pre-composition with h yields a map $(-) \circ h\colon \mathbf{Conv}(X', M) \to \mathbf{Conv}(X, M)$ of effect modules.

In the other direction, for an effect module $E \in \mathbf{EMod}_M$, the homset $\mathbf{EMod}(E, M)$ of effect module maps yields a convex set: for a formal convex sum $\sum_j m_j f_j$, where $f_j\colon E \to M$ in $\mathbf{EMod}_M$ and $m_j \in M$, we can define an actual sum $f\colon E \to M$ by $f(y) = \ovee_j\, m_j \cdot f_j(y)$. This f forms a map of effect modules. Again, functoriality is obtained via pre-composition.

The dual adjunction between $\mathbf{EMod}_M$ and $\mathbf{Conv}_M$ involves a bijective correspondence that is obtained by swapping arguments. $\square$

Finally, via suitable totalisations of the partial operation $\ovee$ (see [36]) one can show that the category $\mathbf{EMod}_{[0,1]}$ is equivalent to the category of (real) partially ordered vector spaces with a unit, see [38].

6 Substitution

In logic, when we have a formula (predicate) $\varphi(y)$ involving a free variable y of type Y, and a term $t\colon X \to Y$, we can perform substitution, and obtain a formula

$$t^*(\varphi) = \varphi[t(x)/y]$$

now involving a free variable x of type X. This substitution operation t^* transforms predicates on Y into predicates on X. It is functorial, in the sense that:

$$\mathrm{id}^*(\varphi) = \varphi \quad \text{and} \quad s^*t^*(\varphi) = (t \circ s)^*(\varphi).$$

In a dynamical setting as described below it makes sense to write these substitution functors as "boxes" in modal/dynamic logic (like in [30]): $[t](\varphi) = t^*(\varphi)$ can be read as the predicate which holds for those states x such that φ holds for $t(x)$. That is, $[t](\varphi)$ says "φ holds after t". Later in Subsection 9.1 we shall see how "tests" can be added.

In the present context where we describe predicates as maps of the form $q\colon Y \to Y + Y$ with $\nabla \circ q = \mathrm{id}$ we can also define substitution. If we have a map $f\colon X \to Y$ we would like to obtain a predicate $f^*(q) = [f](q)$ via substitution. If we assume that in the category where we are working the diagram (D) in Assumption 1 is a pullback, we can obtain a predicate $f^*(q)$ on X in:

$$\begin{tikzcd}
X \arrow[rr, "f"] \arrow[dr, dashed, "f^*(q)"] \arrow[ddr, bend right, equal] & & Y \arrow[dr, bend left, "q"] & \\
& X+X \arrow[rr, "f+f"] \arrow[d, "\nabla"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[drr, phantom, "(D)"] & & Y+Y \arrow[d, "\nabla"] \\
& X \arrow[rr, "f"'] & & Y
\end{tikzcd} \tag{7}$$

Proposition 3. *Let **C** be a category satisfying Assumption 1. The mapping $X \mapsto Pred(X)$ then extends to a functor:*

$$\boldsymbol{C}^{op} \xrightarrow{Pred} \boldsymbol{EA}$$

*where, on an arrow $f\colon X \to Y$ in **C**, one writes $f^* = Pred(f)\colon Pred(Y) \to Pred(Y)$.*

Such a functor is also called an "indexed category"; it forms a common structure in categorical logic. For instance, sending an object X in a category **B** with pullbacks to the poset $Sub(X)$ of subobjects on X, gives rise to a similar functor $\mathbf{B}^{\mathrm{op}} \to \mathbf{PoSets}$.

Proof. Using the uniqueness of the mediating map in pullbacks one easily checks that substitution is functorial: $\mathrm{id}^* = \mathrm{id}$ and $(g \circ f)^* = f^* \circ g^*$. In a similar way one checks that f^* preserves the effect algebra structure. For instance $f^*(1) = 1$ since:

$$(f + f) \circ f^*(1) = 1 \circ f = \kappa_1 \circ f = (f + f) \circ \kappa_1 = (f + f) \circ 1.$$

Now assume $q \perp q'$ in $\mathit{Pred}(Y)$, say via bound $b\colon Y \to (Y+Y)+Y$. Then there is a bound c for $f^*(q)$ and $f^*(q')$, obtained via stacked pullbacks (D) and (D+) from Assumption 1.(iii) in:

$$\begin{tikzcd}
X \arrow[r, "f"] \arrow[dr, dashed, "c"] \arrow[dddr, equal, bend right] & Y \arrow[dr, bend left, "b"] & \\
& (X+X)+X \arrow[r, "(f+f)+f"] \arrow[d, "\nabla+\mathrm{id}"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "(D+)"] & (Y+Y)+Y \arrow[d, "\nabla+\mathrm{id}"] \\
& X+X \arrow[r, "f+f"] \arrow[d, "\nabla"'] \arrow[dr, phantom, "\lrcorner", very near start] \arrow[dr, phantom, "(D)"] & Y+Y \arrow[d, "\nabla"] \\
& X \arrow[r, "f"'] & Y
\end{tikzcd}$$

The outer diagram commutes since $\nabla \circ (\nabla + \mathrm{id}) \circ b \circ f = \nabla \circ (q \ovee q') \circ f = \mathrm{id} \circ f = f$. We obtain $[\mathrm{id}, \kappa_2] \circ c = f^*(q)$ and $[[\kappa_2, \kappa_1], \kappa_2] \circ c = f^*(q')$ via uniqueness. Hence c is a bound for $f^*(q)$ and $f^*(q')$ in $\mathit{Pred}(Y)$. One then uses diagram (D) as above to prove that $f^*(q) \ovee f^*(q') = (\nabla + \mathrm{id}) \circ c$ and $f^*(p \ovee q')$ are equal. $\square$

Assuming this indexed category $\mathit{Pred}\colon \mathbf{C}^{\mathrm{op}} \to \mathbf{EA}$ we shall write $\mathit{Pred}(\mathbf{C})$ for the associated "Grothendieck" completion. This category $\mathit{Pred}(\mathbf{C})$ has predicates $p\colon X \to X+X$ as objects. A morphism $\big(X \overset{p}{\to} X+X\big) \xrightarrow{f} \big(Y \overset{q}{\to} Y+Y\big)$ in $\mathit{Pred}(\mathbf{C})$ is given by a map $f\colon X \to Y$ in **C** between the underlying object that satisfies $p \le f^*(q)$ — where $\le$ is the order in the effect algebra $\mathit{Pred}(X)$ of predicates on X. There is then an obvious forgetful functor:

$$\begin{tikzcd}
\mathit{Pred}(\mathbf{C}) \arrow[d] \\
\mathbf{C}
\end{tikzcd} \tag{8}$$

that is a (split) fibration (see [32]). Sending an object X to falsum $\kappa_2\colon X \to X+X$ or to truth $\kappa_1\colon X \to X+X$ yields both a left and right adjoint (respectively) to this forgetful functor.

In the language of fibrations, the category $\mathit{Pred}(\mathbf{C})$ is called the *total category*, and a poset $\mathit{Pred}(X)$ is called *fibre* or *fibre category* over X.

6.1 Extensive categories

We have already seen in Lemma 1 that an extensive category **C** satisfies Assumption 1. Thus, Proposition 3 applies, so that we have an indexed category $\mathbf{C}^{\mathrm{op}} \to \mathbf{EA}$, and a fibration (8), capturing the logic of the category **C**. This applies for instance for $\mathbf{C} = \mathbf{Sets}$, or any other topos. Substitution (7) corresponds in this case to inverse images: $f^*(p_U) = p_{f^{-1}(U)}$, using the predicate notation p_U from Example 1. And when we identify predicates on Y with maps $Y \to 2 = 1+1$, then substitution is simply precomposition.

Recall, that we describe our logic of predicates in terms of effect algebras, and not in terms of Heyting algebras, as is usually done for a topos. Below we show that this (effect) logic involves “comprehension”, that is, a mapping turning a predicate p on X into a sub-set/type $\{p\} \hookrightarrow X$ of those element satisfying p. In categorical logic, comprehension is described via a right adjoint to the truth functor.

Proposition 4. *The effect algebra logic of an extensive category* $\mathbf{C}$ *has comprehension: the truth functor* $1\colon \mathbf{C} \to \mathit{Pred}(\mathbf{C})$ *has a right adjoint* $\{-\}$*, giving us two successive adjoints to the forgetful functor (fibration) in:*

$$\begin{tikzcd}
\mathit{Pred}(\boldsymbol{C}) \arrow[d, "\dashv"] \arrow[d, bend left=70, "\{-\}"] \\
\boldsymbol{C} \arrow[u, bend right=40, "1"', "\dashv"]
\end{tikzcd}$$

Proof. For a predicate $q\colon Y \to Y+Y$ we define $\{q\} \in \mathbf{C}$ via the pullback:

$$\begin{tikzcd}
\{q\} \arrow[r, tail, "m"] \arrow[d, "q_1"'] \arrow[dr, phantom, "\lrcorner", very near start] & Y \arrow[d, "q"] \\
Y \arrow[r, tail, "\kappa_1"'] & Y+Y
\end{tikzcd} \tag{9}$$

In this pullback we have $m = q_1$, since $q_1 = \nabla \circ \kappa_1 \circ q_1 = \nabla \circ q \circ m = m$. The adjunction $1 \dashv \{-\}$ involves the following bijective correspondence between maps in $\mathit{Pred}(\mathbf{C})$ and in $\mathbf{C}$:

$$\begin{array}{c}
1X = \begin{pmatrix} X \\ \big\downarrow{\scriptstyle \kappa_1} \\ X+X \end{pmatrix} \xrightarrow{\;f\;} \begin{pmatrix} Y \\ \big\downarrow{\scriptstyle q} \\ Y+Y \end{pmatrix} \\
\hline\hline
X \xrightarrow[g]{\quad} \{q\}
\end{array}$$

This works as follows.

- Given a map f in $\mathit{Pred}(\mathbf{C})$ as indicated, then $f\colon X \to Y$ satisfies $1 \le f^*(q)$. This means that $\kappa_1 = 1 = f^*(q)$, and thus $q \circ f = (f+f) \circ q^*(f) = (f+f) \circ \kappa_1 = \kappa_1 \circ f$ in (7). We get a unique map $\overline{f}\colon X \to \{q\}$ via the pullback (9) with $q_1 \circ \overline{f} = f$.
- Given $g\colon X \to \{q\}$ in $\mathbf{C}$, we define $\overline{g} = q_1 \circ g\colon X \to Y$. It is a map of predicates $1X \to q$, since:

$$(\overline{g} + \overline{g}) \circ \kappa_1 = \kappa_1 \circ q_1 \circ g = q \circ m \circ g = q \circ q_1 \circ g = q \circ \overline{g}.$$

Clearly, $\overline{\overline{f}} = f$ and $\overline{\overline{g}} = g$. $\square$

6.2 Dagger biproduct categories

Recall that for an object X in a dagger biproduct category $\mathbf{C}$ we write $\mathit{pPred}(X)$ for the collection of positive predicates on X. If $\mathbf{C}$ is additive and zerosumfree (*i.e.* $f + g = 0 \Rightarrow f = g = 0$ for positive f, g), this is an effect algebra, see Proposition 1. We shall first discuss substitution f^* along unitary maps f, because it can be done via the pullback (7) like in the above extensive case. Once we have derived an explicit formula for this substitution, using the pullback property, we realise that this formula can be applied more generally for dagger monos. This is formulated as a second, more general proposition below.

Proposition 5. *For each zerosumfree dagger additive category* $\boldsymbol{C}$*, on the base category* $\dagger$-*Iso*$(\boldsymbol{C})$ *with unitary maps as morphisms, there is an indexed category of positive predicates,*

$$\dagger\text{-}\mathit{Iso}(\boldsymbol{C})^{op} \xrightarrow{\mathit{pPred}} \boldsymbol{EA}$$

in which substitution along a unitary map $f\colon X \to Y$ *is given by the formula:*

$$f^*(q) \,=\, (f^\dagger \oplus f^\dagger) \circ q \circ f. \tag{10}$$

Proof. If we restrict ourselves to the subcategory $\dagger$-*Iso*$(\mathbf{C}) \hookrightarrow \mathbf{C}$ of unitary maps, we know that diagrams (D) and (D+) are pullbacks by Lemma 2, and so we can define substitution via the pullback (7) in the beginning of this section—and use (D+) to prove that substitution preserves $\ovee$, see the proof of Proposition 3.

The equation $f^*(q) = (f^\dagger \oplus f^\dagger) \circ q \circ f$, for a unitary map f, follows by uniqueness of mediating maps for the pullback (7):

$$\begin{aligned} \nabla \circ (f^\dagger \oplus f^\dagger) \circ q \circ f \,&=\, f^\dagger \circ \nabla \circ q \circ f \,=\, f^\dagger \circ f \,=\, \mathrm{id} \\ (f \oplus f) \circ (f^\dagger \oplus f^\dagger) \circ q \circ f \,&=\, q \circ f. \end{aligned} \qquad \Box$$

The equation (10) is important, since it incorporates the Born rule, see Example 13 below. In the formula (10) for substitution f^* the map f occurs twice, once before the predicate and once afterwards. This double use of f leads to the squaring of amplitudes, in the computation of probabilities in a quantum setting. In Section 9 we have to take square roots in the quantum case to undo this squaring in substitution.

Here we notice that the formulation (10) allows us to describe substitution f^* not only for dagger isomorphisms (that is, unitary maps), but also for dagger monomorphisms (or isometries). We recall that a dagger mono is a map f with $f^\dagger \circ f = \mathrm{id}$. We write $\dagger$-*Mono*$(\mathbf{C}) \hookrightarrow \mathbf{C}$ for the subcategory with all objects and only dagger monos as morphisms. Then we can extend the previous proposition.

Proposition 6. *Each zerosumfree dagger additive category* $\boldsymbol{C}$ *yields an indexed category:*

$$\dagger\text{-}\mathit{Mono}(\boldsymbol{C})^{op} \xrightarrow{\mathit{pPred}} \boldsymbol{EA}$$

in which substitution along a dagger mono $f\colon X \rightarrowtail Y$ *is given by the formula* (10).

Proof. The equation $\nabla \circ f^*(q) = \mathrm{id}$ holds for $f^*(q) = (f^\dagger \oplus f^\dagger) \circ q \circ f$ since f is a dagger mono. Obviously such substitution f^* is functorial. Also, it preserves the effect algebra structure:

$$f^*(1) \,=\, (f^\dagger \oplus f^\dagger) \circ \kappa_1 \circ f \,=\, \kappa_1 \circ f^\dagger \circ f \,=\, \kappa_1 \,=\, 1.$$

If $q, q'\colon Y \to Y \oplus Y$ have bound $b\colon Y \to (Y \oplus Y) \oplus Y$, then $c = ((f^\dagger \oplus f^\dagger) \oplus f^\dagger) \circ b \circ f$ is a bound for $f^*(q)$ and $f^*(q')$ showing $f^*(q \ovee q') = f^*(q) \ovee f^*(q')$. $\Box$

The Born rule that describes the probability of an outcome of a measurement on states is obtained in the current framework as a special instance of the substitution formula (10).

Example 13. Assume an arbitrary qubit $x = (\alpha, \beta) \in \mathbb{C}^2 = \mathbb{C} \oplus \mathbb{C}$, where $|\alpha|^2 + |\beta|^2 = 1$. We can also write it as $x = \alpha|0\rangle + \beta|1\rangle$, where vectors $|0\rangle = \left(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right)$ and $|1\rangle = \left(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}\right)$ form the standard basis of $\mathbb{C}^2$. Yet another way is to write the qubit x as linear map in **FdHilb**:

$$\mathbb{C} \xrightarrow{x = \left(\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right)} \mathbb{C}^2 \qquad \text{with} \qquad \mathbb{C}^2 \xrightarrow{x^\dagger = \left(\begin{smallmatrix} \overline{\alpha} & \overline{\beta} \end{smallmatrix}\right)} \mathbb{C}$$

where $x^\dagger \circ x = \alpha\overline{\alpha} + \beta\overline{\beta} = |\alpha|^2 + |\beta|^2 = 1$. Hence this qubit x is a dagger mono. As a result of Proposition 6 this qubit x yields a substitution operation:

$$\mathit{pPred}(\mathbb{C}^2) \xrightarrow{\quad x^* \quad} \mathit{pPred}(\mathbb{C}) \cong [0, 1]$$

This substitution map x^* gives the probability $x^*(p) \in [0, 1]$ that a predicate $p \in \mathit{pPred}(\mathbb{C}^2)$ holds for the state x. It is essentially the *Born rule*. For instance, if we wish to obtain the probability that state x is $|0\rangle$ we compute x^* on the predicate $p_{|0\rangle}$ associated with the qubit $|0\rangle$. Formally, this predicate is the map $p_{|0\rangle} \colon \mathbb{C}^2 \to \mathbb{C}^2 \oplus \mathbb{C}^2$ given by the tuple:

$$p_{|0\rangle} \;=\; \langle\, |0\rangle\langle 0\,|,\; \mathrm{id} - |0\rangle\langle 0\,|\,\rangle \;=\; \langle\, |0\rangle\langle 0\,|,\; |1\rangle\langle 1\,|\,\rangle \;=\; \langle \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right),\; \left(\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}\right) \rangle.$$

Recall from Example 5 that such a predicate can be identified with the first projection $\pi_1 \circ p_{|0\rangle} = |0\rangle\langle 0\,| \colon \mathbb{C}^2 \to \mathbb{C}^2$, which is an effect. The resulting outcome in $[0, 1] \cong \mathit{pPred}(\mathbb{C})$ of substitution x^* is also obtained via such a first projection:

$$\begin{array}{rcl}
\pi_1 \circ x^*(p_{|0\rangle}) & \overset{(10)}{=} & \pi_1 \circ (x^\dagger \oplus x^\dagger) \circ p_{|0\rangle} \circ x \\
& = & x^\dagger \circ \pi_1 \circ p_{|0\rangle} \circ x \\
& = & x^\dagger |0\rangle\langle 0\,| x \\
& = & \left(\begin{smallmatrix} \overline{\alpha} & \overline{\beta} \end{smallmatrix}\right) \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right) \left(\begin{smallmatrix} \alpha \\ \beta \end{smallmatrix}\right) \;=\; \overline{\alpha}\alpha \;=\; |\alpha|^2.
\end{array}$$

The formula (10) for x^* describes probabilities according to the Born rule. Hence, our conclusion: the Born rule is a special case of substitution in (categorical) logic.

More generally, if we have a positive predicate $p = \langle p_1, p_2 \rangle \colon \mathbb{C}^2 \to \mathbb{C}^2 \oplus \mathbb{C}^2$, where the effect $p_1 \colon \mathbb{C}^2 \to \mathbb{C}^2$ has spectral decomposition:

$$p_1 \;=\; \lambda_1 |v_1\rangle\langle v_1\,| + \lambda_2 |v_2\rangle\langle v_2\,|$$

with eigenvalues $\lambda_i \in [0, 1]$ and corresponding eigenvectors v_i, then v_i is also an eigenvector of $p_2 = \mathrm{id} - p_1$, with eigenvalue $1 - \lambda_i$, so that we can write:

$$p_2 \;=\; (1 - \lambda_1)|v_1\rangle\langle v_1\,| + (1 - \lambda_2)|v_2\rangle\langle v_2\,|.$$

Applying substitution x^* to the effect p_1 gives the (general) Born probability rule:

$$\begin{aligned} x^*(p_1) \;=\; x^\dagger p_1 x &\;=\; \lambda_1 x^\dagger |\, v_1 \,\rangle\langle\, v_1 \,| x + \lambda_2 x^\dagger |\, v_2 \,\rangle\langle\, v_2 \,| x \\ &\;=\; \lambda_1 \overline{\langle v_1 \,|\, x\rangle}\langle v_1 \,|\, x\rangle + \lambda_2 \overline{\langle v_2 \,|\, x\rangle}\langle v_2 \,|\, x\rangle \\ &\;=\; \lambda_1 \big|\langle v_1 \,|\, x\rangle\big|^2 + \lambda_2 \big|\langle v_2 \,|\, x\rangle\big|^2. \end{aligned}$$

Notice that the eigenvalues λ_i of the predicate/effect act as weights (multiplication factors) in this probability formula — as discussed in the introduction.

Also in this dagger biproduct case we can do comprehension, if there is a bit more structure around, namely dagger kernels (see Remark 3).

Proposition 7. *Let* **C** *be a zerosumfree dagger additive category as in Proposition 6, which additionally has dagger kernels. The fibration pPred*$(\mathbf{C}) \rightarrow \dagger$*-Mono*$(\mathbf{C})$ *of positive predicates then has comprehension.*

Proof. The truth functor $1\colon \dagger$*-Mono*$(\mathbf{C}) \rightarrow$ *pPred*$(\mathbf{C})$ sends an object X to the truth predicate $\kappa_1\colon X \rightarrow X \oplus X$. In the other direction the comprehension functor $\{-\}$ sends a predicate $q = \langle q_1, q_2\rangle\colon Y \rightarrow Y \oplus Y$ to (the domain of) the kernel of the second component:

$$\{q\} \stackrel{\text{def}}{=} \ker(q_2) \;=\; \ker(\mathrm{id} - q_1) \;=\; \mathrm{eq}(\mathrm{id}, q_1),$$

where eq is written for the equaliser. We briefly check the adjunction $1 \dashv \{-\}$. We shall write the kernel map explicitly as $\pi_q\colon \{q\} \rightarrowtail X$, which is a dagger mono.

Assume a dagger mono $f\colon X \rightarrowtail Y$ with $1 \leq f^*(q)$. This means $f^\dagger \circ q_1 \circ f = \mathrm{id}$ and $f^\dagger \circ q_2 \circ f = 0$. Since $f^\dagger$ is epic, we may conclude $q_2 \circ f = 0$. Hence there is a unique map $\overline{f}\colon X \rightarrow \{q\}$ with $\pi_q \circ \overline{f} = f$.

Conversely, assume we have a dagger mono $g\colon X \rightarrowtail \{q\}$. Write $\overline{g} = \pi_q \circ g$. Then:

$$\begin{aligned} \pi_1 \circ \overline{g}^*(q) \;=\; \overline{g}^\dagger \circ q_1 \circ \overline{g} &\;=\; g^\dagger \circ (\pi_q)^\dagger \circ q_1 \circ \pi_q \circ g \\ &\;=\; g^\dagger \circ (\pi_q)^\dagger \circ \pi_q \circ g \qquad \text{since } \pi_q \text{ equalises } \mathrm{id}, q_1 \\ &\;=\; \mathrm{id} \\ \pi_2 \circ \overline{g}^*(q) \;=\; \overline{g}^\dagger \circ q_2 \circ \overline{g} &\;=\; g^\dagger \circ (\pi_q)^\dagger \circ q_2 \circ \pi_q \circ g \\ &\;=\; g^\dagger \circ (\pi_q)^\dagger \circ 0 \circ g \\ &\;=\; 0. \end{aligned}$$

Hence $\overline{g}^*(q) = \langle \mathrm{id}, 0\rangle = \kappa_1 = 1$. Thus $\overline{g}$ is a map $1(X) \rightarrow q$ in *pPred*$(\mathbf{C})$. □

Example 14. Propositions 6 also applies to the category **BifMRel** of sets and bifinite relations. Recall from Example 4 that the positive predicates on X in **BifMRel** are positive matrices $M \in \mathbb{C}^{X\times X}$ with entries in $[0,1]$ on its diagonal. A dagger mono $r\colon X \rightarrowtail Y$ gives rise to a substitution functor $r^*\colon [0,1]^{Y\times Y} \rightarrow [0,1]^{X\times X}$ given by:

$$r^*(N)(x, x') \;=\; \textstyle\sum_{y,y'} r(x,y) \cdot N(y,y') \cdot \overline{r(x',y')}. \tag{11}$$

This equation follows from (10): for a predicate $q\colon Y \to Y+Y$ representing matrix $N \in [0,1]^{Y\times Y}$ as $N(y,y') = q(y,\kappa_1 y')$, we get as resulting matrix in $[0,1]^{X\times X}$,

$$\begin{aligned}
r^*(q)(x,\kappa_1 x') &\overset{(10)}{=} \big((r^\dagger + r^\dagger)\circ q \circ r\big)(x,\kappa_1 x') \\
&= \textstyle\sum_{y\in Y, u\in Y+Y} r(x,y)\cdot q(y,u)\cdot (r^\dagger + r^\dagger)(u,\kappa_1 x') \\
&= \textstyle\sum_{y,y'\in Y} r(x,y)\cdot q(y,\kappa_1 y')\cdot (r^\dagger + r^\dagger)(\kappa_1 y',\kappa_1 x') \\
&= \textstyle\sum_{y,y'\in Y} r(x,y)\cdot N(y,y')\cdot q^\dagger(y',x') \\
&= \textstyle\sum_{y,y'\in Y} r(x,y)\cdot N(y,y')\cdot \overline{q(x',y')}.
\end{aligned}$$

Remark 4. As illustrated above, in dagger categories one can perform substitution f^* for dagger monos f, and not for all maps. This may look like a restriction. But in Hilbert spaces dagger monos contain all the computationally relevant information: as a consequence of Stinespring's (dilation) theorem, each completely positive trace preserving operator (channel) F can be written as $F(A) = f^\dagger \circ (A\otimes \mathrm{id}) \circ f = f^*(A \otimes \mathrm{id})$, for some dagger mono $f\colon H \rightarrowtail H\otimes K$, where K is sometimes called the "ancilla" system, see [29].

6.3 Sets and discrete probability distributions

In Example 3 we have seen that in the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the (discrete probability) distribution monad $\mathcal{D}$ predicates $X \to X+X$ on X correspond to fuzzy predicates $X \to [0,1]$, that is, *Pred*$(X) \cong [0,1]^X$. The question we would like to address is if this also gives rise to an indexed category *Pred*: $\mathcal{K}\ell(\mathcal{D})^{\mathrm{op}} \to$ **EA**.

The diagrams (D) and (D+) from Assumption 1 are not pullbacks in $\mathcal{K}\ell(\mathcal{D})$. Hence the approach followed above, via universality in (7), does not work. But we can proceed in the spirit of this approach, like we did for dagger monos in the previous subsection.

So assume a map $f\colon X \to Y$ in $\mathcal{K}\ell(D)$, describing a stochastic matrix. We would like to have a map of effect algebras $f^*\colon [0,1]^Y \to [0,1]^X$. There is an obvious way to do so via a (finite) sum:

$$f^*(\psi)(x) = \sum_{y\in Y} f(x)(y)\cdot\psi(y). \tag{12}$$

(This sum is finite because by definition a probability distribution $f(x) \in \mathcal{D}(Y)$ has finite support.)

If we understand the fuzzy predicate $\psi \in [0,1]^Y$ as a "weighted event" $E \subseteq Y$, given by the subset $E = \{y \mid \psi(y) \neq 0\} \subseteq Y$, where for $y \in E$ the factor $\psi(y) \in (0,1]$ describes a weight, then $f^*(\psi)$ describes translation of the weighted event ψ on Y to a weighted event on X, with weights as described above in (12).

We get $f^*(1) = 1$, since $f(x) \in \mathcal{D}(Y)$; also $\ovee$ is preserved by substitution. Notice that 'probabilistic' substitution f^* in (12) does not involve a double occurrence of the map f, like in the quantum case (10).

The following result shows that in the probabilistic case one has the same categorical structure as before.

Proposition 8. *There is an indexed category* $\mathit{Pred}\colon \mathcal{K}\ell(\mathcal{D})^{op} \to \boldsymbol{EA}$. *The resulting fibration,*

$$\begin{tikzcd}
\mathit{Pred}(\mathcal{K}\ell(\mathcal{D})) \arrow[d] \\
\mathcal{K}\ell(\mathcal{D})
\end{tikzcd}$$

has a truth functor $1\colon \mathcal{K}\ell(\mathcal{D}) \to \mathit{Pred}(\mathcal{K}\ell(\mathcal{D}))$ *and comprehension* $\{-\}\colon \mathit{Pred}(\mathcal{K}\ell(\mathcal{D})) \to \mathcal{K}\ell(\mathcal{D})$ *as its right adjoint.*

Proof. We indeed have a truth functor $1\colon \mathcal{K}\ell(\mathcal{D}) \to \mathit{Pred}(\mathcal{K}\ell(\mathcal{D}))$ since $f^*(1) = 1$. Its right adjoint $\{-\}$ is defined on $\psi\colon Y \to [0,1]$ as:

$$\{\psi\} \;=\; \{y \in Y \mid \psi(y) = 1\}.$$

We have an inclusion map $\pi_\psi\colon \{\psi\} \hookrightarrow Y$ in **Sets**, yielding a map $\mathcal{J}(\pi_\psi) = \eta \circ \pi_\psi\colon \{\psi\} \to Y$ in $\mathcal{K}\ell(\mathcal{D})$.

We briefly check the adjunction $1 \dashv \{-\}$. Suppose we have a map $f\colon 1(X) \to q$ in *Pred*$(\mathcal{K}\ell(\mathcal{D})$. This means $1 \le \sum_y f(x)(y)\cdot\psi(y)$, for each $x \in X$. Since $\sum_y f(x)(y) = 1$, this can only be the case if $\psi(y) = 1$ when $f(x)(y) \neq 0$. Thus *supp*$(f(x)) \subseteq \{\psi\}$, so that we get a function $\overline{f}\colon X \to \mathcal{D}(\{q\})$ with $\mathcal{D}(\pi_\psi) \circ \overline{f} = f$. In $\mathcal{K}\ell(\mathcal{D})$ this means $\mathcal{J}(\pi_\psi) \circ \overline{f} = f$.

In the other direction, assume we have a map $g\colon X \to \{q\}$ in $\mathcal{K}\ell(\mathcal{D})$. For each $x \in X$ we have $g(x) \in \mathcal{D}(\{\psi\})$. We formally define a function $\overline{g} = \mathcal{D}(\pi_\psi) \circ g\colon X \to \mathcal{D}(Y)$. It yields:

$$\begin{aligned}
1 \;\le\; 1 \;=\; \textstyle\sum_{y\in\{\psi\}} g(x)(y) \;&=\; \textstyle\sum_{y\in\{\psi\}} g(x)(y)\cdot\psi(y) \\
&=\; \textstyle\sum_{y\in Y} \overline{g}(x)(y)\cdot\psi(y) \;=\; \overline{g}^*(\psi)(x).
\end{aligned}$$

Hence $\overline{g}$ is a map $1(X) \to \psi$ in the category *Pred*$(\mathcal{K}\ell(\mathcal{D}))$. $\square$

It is not hard to see that the standard embedding $\mathcal{J}\colon \mathbf{Sets} \to \mathcal{K}\ell(\mathcal{D})$ lifts to a fibred functor:

$$\begin{tikzcd}
\mathit{Pred}(\mathbf{Sets}) \arrow[r] \arrow[d] & \mathit{Pred}(\mathcal{K}\ell(\mathcal{D})) \arrow[d] \\
\mathbf{Sets} \arrow[r, "\mathcal{J}"] & \mathcal{K}\ell(\mathcal{D})
\end{tikzcd}$$

Along the same lines, one might expect a functor $\mathcal{Q}\colon \dagger$-*Mono*(**BifMRel**) $\to \mathcal{K}\ell(\mathcal{D})$, namely: $\mathcal{Q}(X) = X$ on an object X, and $\mathcal{Q}(r)\colon X \to \mathcal{D}(Y)$, for a dagger mono $r\colon X \to Y$, given by:

$$\mathcal{Q}(r)(x)(y) \;=\; \big|r(x,y)\big|^2.$$

This $\mathcal{Q}(r)$ yields a distribution, but we do *not* get a functor since $\mathcal{Q}$ does not preserve composition.

At the end of this section we see that for classical, probabilistic, and quantum logic we have similar fibred categories:

$$\begin{tikzcd}
\mathit{Pred}(\mathbf{Sets}) \arrow[d] & \mathit{Pred}(\mathcal{K}\ell(\mathcal{D})) \arrow[d] & \mathit{pPred}(\mathbf{BifMRel}) \arrow[d] & \mathit{pPred}(\mathbf{Hilb}) \arrow[d] \\
\mathbf{Sets} & \mathcal{K}\ell(\mathcal{D}) & \dagger\text{-}\mathit{Mono}(\mathbf{BifMRel}) & \dagger\text{-}\mathit{Mono}(\mathbf{Hilb})
\end{tikzcd} \tag{13}$$

starting from predicates of the form $X \to X + X$. These four fibrations are fibred effect algebras: each fibre is an effect algebra, and substitution functors preserve the effect algebra structure. The sum $\ovee$ in these fibres is defined in a uniform manner, via bounds as in Definition 3. The substitution functors f^* do not follow the same pattern in all four cases. However, the set-theoretic f^* defined via (7) and quantum f^* via (10) for unitary f both arise via the same pullback.

All the fibrations/logics (13) have the adjunction "truth $\dashv$ comprehension" in common. We conclude this section by showing that the special predicate Ω from Example 6 gives in all these cases the same structure.

Lemma 10. *For all four fibrations in* (13) *the Ω predicate from Example 6 yields a fibred functor $X \mapsto \Omega_X$ from the base category to the total category.*

Proof. Recall, for each object X in the base category we have a predicate $\Omega_X = \kappa_1 + \kappa_2 \colon X + X \to (X+X)+(X+X)$. On a map $f \colon X \to Y$ one gets $f + f \colon \Omega_X \to \Omega_Y$. This is a cartesian map since for each of the four fibrations one has $(f+f)^*(\Omega_Y) = \Omega_X$. □

7 Effect monoids of predicates

This section shows how predicates carry a (multiplicative) monoid structure $\cdot$ that captures both intersection for subsets and (pointwise) multiplication for fuzzy predicates. This total multiplication operation may seem more natural than the partial sum $\ovee$, but it is not so well-behaved: it is not always preserved by substitution and its definition does not work for positive predicates. As we shall see, this multiplication only works well on the tensor unit — if any. Thus, it will be used especially for probabilities, *i.e.* for predicates on a tensor unit.

Assume a category **C** with coproducts. The set of predicates $\mathit{Pred}(X)$ on an object X carries a monoid structure which we shall write as multiplication. For two predicates $p, q \colon X \to X + X$ define:

$$p \cdot q = \Big(X \xrightarrow{q} X + X \xrightarrow{p+\mathrm{id}} (X+X)+X \xrightarrow{[\mathrm{id},\kappa_2]} X + X \Big). \tag{14}$$

That is, $p \cdot q = [p, \kappa_2] \circ q$.

Lemma 11. *Consider multiplication* (14) *of predicates on X, as described above.*

(i). It yields a monoid structure on $\mathit{Pred}(X)$ with the truth predicate 1 *as unit element.*

(ii). In case Pred(X) is an effect algebra, via sum $\ovee$ as in Definition 3, then multiplication is a bihomomorphism, making each Pred(X) an effect monoid.

(iii). Multiplication is preserved by substitution functors f^*, as defined in the pullback (7).

Proof. (i). We check associativity:

$$\begin{aligned} p \cdot (q \cdot r) \;&=\; [p, \kappa_2] \circ (q \cdot r) \;=\; [p, \kappa_2] \circ [q, \kappa_2] \circ r \\ &=\; [[p, \kappa_2] \circ q, \kappa_2] \circ r \;=\; [p \cdot q, \kappa_2] \circ r \;=\; (p \cdot q) \cdot r. \end{aligned}$$

It is similarly easy to see that $1 = \kappa_1$ is neutral element for this multiplication:

$$p \cdot 1 \;=\; [p, \kappa_2] \circ \kappa_1 \;=\; p \qquad\qquad 1 \cdot q \;=\; [\kappa_1, \kappa_2] \circ q \;=\; q.$$

(ii). In the same way one checks that $0 \cdot p = 0 = p \cdot 0$. We concentrate on $\ovee$. Let $q, q' \colon X \to X + X$ be orthogonal predicates, say via bound $b \colon X \to (X + X) + X$. We define the following two new bounds.

$$c \;=\; \left(\begin{tikzcd} X \arrow[d, "p"] \\ X + X \arrow[d, "b + \mathrm{id}"] \\ ((X + X) + X) + X \arrow[d, "{[\mathrm{id}, \kappa_2]}"] \\ (X + X) + X \end{tikzcd}\right) \qquad\qquad d \;=\; \left(\begin{tikzcd} X \arrow[d, "b"] \\ (X + X) + X \arrow[d, "(p + p) + \mathrm{id}"] \\ ((X + X) + (X + X)) + X \arrow[d, "{[[\kappa_1 + \mathrm{id}, \kappa_2 + \mathrm{id}], \kappa_2]}"] \\ (X + X) + X \end{tikzcd}\right)$$

The bound c shows $q \cdot p \perp q' \cdot p$ and $(q \cdot p) \ovee (q' \cdot p) = (q \ovee q') \cdot p$. Similarly, d proves $p \cdot q \perp p \cdot q'$ and $(p \cdot q) \ovee (p \cdot q') = p \cdot (q \ovee q')$.

(iii). For a map $f \colon X \to Y$ and predicates $p, q \colon Y \to Y + Y$ we have $f^*(p \cdot q) = f^*(p) \cdot f^*(q)$ by an easy uniqueness argument. $\square$

With this result we can strengthen Proposition 3.

Corollary 1. *For each category **C** satisfying Assumption 1 there is a functor (indexed category):*

$$\mathbf{C}^{op} \xrightarrow{\;\;Pred\;\;} \mathbf{EMon}$$

Example 15. In the category **Sets** we have seen in Example 1 that subsets $U \subseteq X$ correspond to predicates $p_U \colon X \to X + X$, given by: $p_U(x) = \kappa_1 x$ iff $x \in U$. It is not hard to check that $p_U \cdot p_V = p_{U \cap V}$. Hence in this case the multiplicative structure on predicates corresponds to intersection.

In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of sets and stochastic relations predicates on X correspond to fuzzy predicates $\varphi \colon X \to [0, 1]$, via $p_\varphi(x) = \varphi(x)\kappa_1 x + (1 - \varphi(x))\kappa_2 x$. Multiplication $p_\varphi \cdot p_\psi$ describes the predicate $p_{\psi \cdot \varphi}$, where $\psi \cdot \varphi \colon X \to [0, 1]$ is the pointwise multiplication of probabilities.

However, in this case we do *not* get an indexed category *Pred*: $\mathcal{K}\ell(\mathcal{D})^{\mathrm{op}} \to$ **EMon**, since substitution f^* as defined in (12) does not preserve multiplication.

Positive maps between Hilbert spaces are in general not closed under composition. Therefore, effect algebras *pPred*(X) of *positive* predicates do not have a multiplication (14) as used above.

However, in case our dagger biproduct category is also a dagger monoidal category, with tensor $(\otimes, I)$, the positive predicates *pPred*(I) on the tensor unit I do carry a multiplication (14). The reason is that positive maps $I \to I$ on I are closed under composition, essentially because scalars always commute (multiplicatively), see (see [41,2]). Here we need a slightly more general result. It applies to so-called *dagger monoidal* categories, where $(f \otimes g)^\dagger = f^\dagger \otimes g^\dagger$ and all monoidal isomorphisms (see *e.g.* [48, Ch. VII]) are dagger isomorphisms.

Lemma 12. *Let **C** be a dagger monoidal category. If both $s\colon I \to I$ and $t\colon X \to X$ are positive maps, then so is the "product"*

$$s \star x = \Big(X \xrightarrow[\cong]{\lambda^{-1}} I \otimes X \xrightarrow{s \otimes t} I \otimes X \xrightarrow[\cong]{\lambda} X\Big)$$

This operation $\star$ is sometimes called scalar multiplication. Later, in Section 8 we shall describe an analogous operation that we call probability multiplication.

Proof. Since s, t are positive we can write them as $s = f^\dagger \circ f$ and $t = g^\dagger \circ g$, for some maps $f\colon I \to A$ and $g\colon X \to B$. Then $s \star t = h^\dagger \circ h$ for $h = (f \otimes g) \circ \lambda^{-1}\colon I \overset{\cong}{\to} I \otimes I \to A \otimes B$, because $\lambda^{-1} = \lambda^\dagger$. □

The (positive) predicates on the tensor unit will be called probabilities. They do carry a multiplication operation.

Lemma 13. *In a zerosumfree dagger monoidal additive category **C**, the collection pPred(I) of predicates on the tensor unit is an effect monoid, with multiplication $\cdot$ from* (14) *given by composition:*

$$\pi_1 \circ (p \cdot q) \;=\; (\pi_1 \circ p) \circ (\pi_1 \circ q).$$

In Theorem 2 it will be shown that this operation is commutative.

Proof. Since:

$$\begin{aligned}\pi_1 \circ (p \cdot q) \;=\; \pi_1 \circ [p, \kappa_2] \circ q \;&=\; [\pi_1 \circ p, \pi_1 \circ \kappa_2] \circ q \\ &=\; [\pi_1 \circ p, 0] \circ q \\ &=\; \pi_1 \circ p \circ [\mathrm{id}, 0] \circ q \;=\; \pi_1 \circ p \circ \pi_1 \circ q. \qquad \square\end{aligned}$$

8 Effect modules

In this section it will be shown that predicates carry the structure of an effect module: predicates can be multiplied with a probability (scalar). Unlike multiplication of predicates, this probability multiplication is well-behaved.

At this stage we assume our category **C**, which standardly has coproducts $(0, +)$, also has a symmetric monoidal structure $(I, \otimes)$, in such a way that tensors distribute over coproducts. The latter means that the canonical maps:

$$(X \otimes Z) + (Y \otimes Z) \xrightarrow{[\kappa_1 \otimes \mathrm{id}, \kappa_2 \otimes \mathrm{id}]} (X + Y) \otimes Z \qquad\qquad 0 \xrightarrow{\;!\;} 0 \otimes Z$$

are isomorphisms. We shall write *dis*: $(X + Y) \otimes Z \xrightarrow{\cong} (X \otimes Z) + (Y \otimes Z)$ for the inverse of the first map. There is an analogous map *dis*$'$: $Z \otimes (X + Y) \xrightarrow{\cong} (Z \otimes X) + (Z \otimes Y)$.

In case **C** is a dagger biproduct category we also assume such a (distributive) symmetric monoidal structure, where $(I, \otimes)$ is a so-called *dagger monoidal* structure (see before Lemma 12). One can now describe the map *dis* explicitly as *dis* $= \langle \pi_1 \otimes \mathrm{id}, \pi_2 \otimes \mathrm{id} \rangle$.

From the previous section we know that the predicates *Pred*(I) and positive predicates *pPred*(I) on a tensor unit $I \in$ **C** carry a multiplication operation $\cdot$ giving the structure of an effect monoid — assuming additional requirements guaranteeing effect algebra structure, see Section 5. We have described the (positive) predicates on $I \in$ **C** as *probabilities* and write them as Pr(**C**), see Definition 2. We show that predicates (on arbitrary objects) come with a form of scalar multiplication, namely multiplication with probabilities. Such structures are the effect modules from Subsection 5.3.

Definition 6. *Let **C** be a monoidal category with coproducts, where tensors distribute over coproducts. For a probability* $s\colon I \to I + I$ *and a predicate* $p\colon X \to X + X$ *we define the* probability product $s \bullet p\colon X \to X + X$ *as the following composite.*

$$\begin{tikzcd}
X \arrow[r, "p"] & X + X \arrow[d, "\lambda^{-1}+\mathrm{id}"', "\cong"] & & \\
& I \otimes X + X \arrow[r, "s \otimes \mathrm{id} + \mathrm{id}"] & (I + I) \otimes X + X \arrow[d, "dis+\mathrm{id}"', "\cong"] & \\
& & (I \otimes X + I \otimes X) + X \arrow[r, "{[\lambda+\lambda, \kappa_2]}"] & X + X
\end{tikzcd} \tag{15}$$

Before investigating the properties of this scalar multiplication, we briefly look at an example that makes clear what happens in this definition.

Example 16. The Kleisli category $\mathcal{K}\ell(\mathcal{D})$ has monoidal structure $(1, \times)$, essentially because distribution $\mathcal{D}$ is a so-called commutative (or monoidal) monad. These products $\times$ distribute over coproducts, like in **Sets**. Notice that in the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ the set of scalars $1 \to 1$ is a singleton, since the distribution monad is affine: $\mathcal{D}(1) \cong 1$. The probabilities, however, are non-trivial, and correspond to elements of the unit interval $[0, 1]$.

Indeed, a probability $s\colon 1 \to \mathcal{D}(1 + 1)$ can be identified with a fuzzy predicate $1 \to [0, 1]$, and thus with an ordinary probability $s \in [0, 1]$. Next, assume we have an arbitrary predicate $p_\varphi\colon X \to X + X$, determined by a fuzzy predicate $\varphi\colon X \to [0, 1]$ like in Example 3. The multiplication $s \bullet p_\varphi\colon X \to \mathcal{D}(X + X)$ from (15) works as follows.

a. Applying the predicate p to an element x yields a convex sum $p(x) = \varphi(x)\kappa_1 x + (1 - \varphi(x))\kappa_2 x$;

b. The probability s formally is a convex sum $s\kappa_1 * + (1-s)\kappa_2 *$, and turns $p(x)$ into the convex sum:

$$s \cdot \varphi(x)\kappa_1\kappa_1 x + (1-s)\varphi(x)\kappa_2\kappa_1 x + (1 - \varphi(x))\kappa_2 x$$

c. The last two elements of this sum are combined in the outcome:

$$\begin{aligned} & s \cdot \varphi(x)\kappa_1 x + \big((1-s)\varphi(x) + (1-\varphi(x))\big)\kappa_2 x \\ = \; & s \cdot \varphi(x)\kappa_1 x + (1 - s \cdot \varphi(x))\kappa_2 x. \end{aligned}$$

Thus, $s \bullet p_\varphi = p_{s \cdot \varphi}$, where $s \cdot \varphi \colon X \to [0,1]$ describes pointwise multiplication with $s \in [0,1]$. Clearly, such multiplication $\bullet$ is preserved under substitution.

Proposition 9. *Assuming the relevant structure exists, probability multiplication $\bullet$ defined in* (15),

(i). turns sets of predicates Pred(X) and pPred(X) into effect modules;

(ii). is preserved by substitution f^*, as defined in the pullback (7)*;*

(iii). coincides with multiplication $\cdot$ from (14) *for probabilities, i.e. for predicates on the tensor unit I.*

Proof. We leave it to the reader to check that $s \bullet p \colon X \to X + X$ is well-defined, that is, satisfies $\nabla \circ (s \bullet p) = \mathrm{id}$, and also that it is positive when both s, p are positive (via Lemma 12). This $\bullet$ is a bihomomorphism of effect algebras. We prove preservation of $\ovee$ only in the second argument, and leave the first argument to the reader. So assume we have orthogonal predicates $p, q \colon X \to X + X$ with bound $b \colon X \to (X+X)+X$. Given a probability $s \colon I \to I + I$ we define a new bound c as:

$$\begin{tikzcd}
X \arrow[r, "b"] & (X+X)+X \arrow[d, "(\lambda^{-1}+\lambda^{-1})+\mathrm{id}"', "\cong"] & \\
& (I \otimes X + I \otimes X) + X \arrow[r, "(s\otimes\mathrm{id}+s\otimes\mathrm{id})+\mathrm{id}"] & ((I+I)\otimes X + (I+I)\otimes X) + X \arrow[d, "(\mathit{dis}+\mathit{dis})+\mathrm{id}"', "\cong"] \\
& & ((I \otimes X + I \otimes X) + (I \otimes X + I \otimes X)) + X \arrow[d, "{[[(\kappa_1\circ\lambda)+\lambda,(\kappa_2\circ\lambda)+\lambda],\kappa_2]}"'] \\
& & (X+X)+X
\end{tikzcd}$$

This c is a bound for $s \bullet p$ and $s \bullet q$, showing that $s \bullet (p \ovee q) = (s \bullet p) \ovee (s \bullet q)$.

The next point is showing that $\bullet$ is an action, that is, $1 \bullet p = p$ and $(s \cdot t) \bullet p = s \bullet (t \bullet p)$. This is laborious but straightforward.

Preservation of probability multiplication under substitution, that is, $f^*(s \bullet p) = s \bullet f^*(p)$, follows from a simple uniqueness argument in (7).

Finally, for probabilities $s, t \colon I \to I + I$ we use that $\lambda = \rho \colon I \otimes I \xrightarrow{\cong} I$, in order to show that probability multiplication $\bullet$ in (15) is the same as multiplication $\cdot$ in (14):

$$\begin{aligned} s \bullet t \;&=\; [\lambda + \lambda, \kappa_2] \circ (\mathit{dis} + \mathrm{id}) \circ ((s \otimes \mathrm{id}) + \mathrm{id}) \circ (\lambda^{-1} + \mathrm{id}) \circ t \\ &=\; [(\rho + \rho) \circ \mathit{dis} \circ (s \otimes \mathrm{id}) \circ \rho^{-1}, \kappa_2] \circ t \\ &=\; [\rho \circ (s \otimes \mathrm{id}) \circ \rho^{-1}, \kappa_2] \circ t \;=\; [s, \kappa_2] \circ t \;=\; s \cdot t. \end{aligned}$$ □

There is an analogue of the famous Kelly-LaPlaza result (see [41,2]) that states that composition of scalars — *i.e.* of endomaps $I \to I$ on the tensor unit — is commutative. The proof is essentially as in [35].

Theorem 2. *Multiplication of probabilities is commutative.*

Proof. One proves that for two arbitrary probabilities $s, t \colon I \rightrightarrows I + I$ the two products $s \cdot t$ and $t \cdot s$ are both equal to the following composite.

$$\begin{tikzcd}
I \arrow[r, "\lambda^{-1}=\rho^{-1}", "\cong"'] & I \otimes I \arrow[r, "s\otimes t"] & (I+I)\otimes(I+I) \arrow[d, "dis", "\cong"'] & \\
& & \big(I\otimes(I+I)\big) + \big(I\otimes(I+I)\big) \arrow[d, "dis'+dis'", "\cong"'] & \\
& & \big((I\otimes I)+(I\otimes I)\big) + \big((I\otimes I)+(I\otimes I)\big) \arrow[r, "(\rho+\rho)+(\rho\circ\nabla)"] & I+I
\end{tikzcd}$$

where *dis*$'$ is the obvious (swapped) distribution isomorphism. □

Recall that we write $\mathrm{Pr}(\mathbf{C})$ for the collection of probabilities in a category **C** — assuming the relevant structure exists to consider (positive) predicates on the tensor unit $I \in \mathbf{C}$. These probabilities form a commutative effect monoid. The category of effect modules over this monoid is written as $\mathbf{EMod}_{\mathrm{Pr}(\mathbf{C})}$.

The next result is a direct consequence of Proposition 9.

Proposition 10. *For each category **C** satisfying Assumption 1 there is a functor (indexed category):*

$$\mathbf{C}^{op} \xrightarrow{\;Pred\;} \mathbf{EMod}_{Pr(\mathbf{C})}$$ □

If we instantiate this result for **Sets**, we get $\mathrm{Pr}(\mathbf{Sets}) = 2 = \{0,1\}$ and $\mathbf{EMod}_2 = \mathbf{EA}$, since probability multiplication is trivial with truth values as probabilities. In fact, the predicates in this set-theoretic case are not just effect algebras but Boolean algebras.

Proposition 11. *For a zerosumfree dagger monoidal additive category **B** there are indexed categories:*

$$\dagger\text{-}Iso(\mathbf{C})^{op} \xrightarrow{pPred} \mathbf{EMod}_{Pr(\mathbf{C})} \qquad \textit{and} \qquad \dagger\text{-}Mono(\mathbf{C})^{op} \xrightarrow{pPred} \mathbf{EMod}_{Pr(\mathbf{C})}$$

Proof. What remains is to check that substitution (10) preserves probability multiplication. For a dagger mono (or unitary) map $f \colon X \rightarrowtail Y$, a predicate $q \colon Y \to Y \oplus Y$ and a probability s we can prove $f^*(s \bullet q) = s \bullet f^*(q)$ via a direct calculation. □

Example 17. When we instantiate the last result with the category **Hilb** of Hilbert spaces we obtain an indexed category of "effects"

$$\dagger\text{-}Mono(\mathbf{Hilb})^{\mathrm{op}} \xrightarrow{\;\text{effects}\;} \mathbf{EMod}_{[0,1]}$$

As is well-known, the effects form an effect module over the effect monoid $[0,1]$ of ordinary probabilities, see [38,53].

In a similar manner, the category $\mathcal{K}\ell(\mathcal{D})$ of sets and stochastic relations forms the basis for an indexed category:

$$\mathcal{K}\ell(\mathcal{D})^{\mathrm{op}} \xrightarrow{\mathit{Pred}=[0,1]^{(-)}} \mathbf{EMod}_{[0,1]}$$

of fuzzy predicates. Also these fuzzy predicates $[0,1]^X$ form well-known examples of effect modules (over $[0,1]$). What is new here is not only the functoriality of this construction, but also the uniformity, via predicates $X \to X + X$.

9 Characteristic maps

This section introduces characteristic maps char_p for predicates p. These maps will be used for test operations $[p?]$ and $\langle p? \rangle$ in the present section, and for measurement in the next section. Recall the predicate $\Omega_X = \kappa_1 + \kappa_2 \colon X + X \to (X + X) + (X + X)$ from Example 6.

Definition 7. *Let $p \colon X \to X + X$ be a predicate, in a fibred category that supports predicates and substitution. A* characteristic *map for p is a morphism $\mathit{char}_p \colon X \to X + X$ in the base category with:*

$$p = \big(\mathit{char}_p\big)^*(\Omega_X).$$

We intentionally use the notation Ω_X that is standardly used for a subobject classifier in a topos (see *e.g.* [49]). It has the property that for each mono $m \colon U \rightarrowtail X$ there is a unique map $\mathit{char}_U \colon X \to \Omega$ forming a pullback:

$$\begin{tikzcd} U \arrow[r] \arrow[d, tail, "m"'] \arrow[dr, phantom, "\lrcorner", very near start] & 1 \arrow[d, "\top"] \\ X \arrow[r, "\mathit{char}_U"'] & \Omega \end{tikzcd} \qquad \text{saying} \qquad m = \big(\mathit{char}_m\big)^{-1}(\top).$$

Notice that in a topos there is a fixed classifier, whereas in the current situation our $\Omega_X \colon X + X \to (X + X) + (X + X)$ is parametrised by the object X. Also, in a topos characteristic maps are unique. In the present setting there is no uniqueness, but in our main examples we do have a canonical choice for char_p.

It turns out that in our examples of classical, probabilistic and quantum logic such characteristic maps exist. The quantum case is the most interesting.

Lemma 14. *Both in classical and probabilistic predicate logic, of the categories* **Sets** *and of $\mathcal{K}\ell(\mathcal{D})$ respectively, one can take $\mathit{char}_p = p$.*

Proof. In **Sets**, assume $p = p_U \colon X \to X + X$ for $U \subseteq X$ like in Example 1. As shown in Example 6 the predicate Ω_X in **Sets** is the left inclusion $\kappa_1 \colon X \hookrightarrow X + X$. Then:

$$p^*(\Omega_X) = p^{-1}(\kappa_1) = \{x \mid p(x) = \kappa_1 x\} = U.$$

In the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ assume $p = p_\varphi$ for a fuzzy predicate $\varphi \in [0,1]^X$, see Example 3. The predicate Ω_X corresponds to the fuzzy predicate $\psi \in [0,1]^{X+X}$ with $\psi(\kappa_1 x) = 1$ and $\psi(\kappa_2 x) = 0$. Thus:

$$p^*(\Omega_X)(x) \;=\; \textstyle\sum_{u \in X+X} p(x)(u) \cdot \psi(u) \;=\; p(x)(\kappa_1 x) \;=\; \varphi(x). \qquad \square$$

In the set-theoretic case it is easy to see that characteristic maps are not unique. Assume a predicate $p = p_U \colon X \to X + X$ representing $U \subseteq X$. If $f \colon X \to X$ is an isomorphism with $f(x) \in U$ for $x \in U$, then $(f + \mathrm{id}) \circ p \colon X \to X + X$ can also be used as characteristic map for p.

We turn to quantum logic. For a positive predicate $p = \langle p_1, p_2 \rangle \colon X \to X \oplus X$ in the category **FdHilb** of finite-dimensional Hilbert spaces write $\sqrt{p} = \langle \sqrt{p_1}, \sqrt{p_2} \rangle \colon X \to X \oplus X$. The square root $\sqrt{f}$ of a positive endomap $f \colon X \to X$ in **FdHilb** is obtained as operator function (see *e.g.* [51, §§2.1.8]): write f as spectral decomposition $f = \sum_i \lambda_i |\, v_i \,\rangle\langle\, v_i \,|$, and define $\sqrt{f} = \sum_i \sqrt{\lambda_i} |v_i\rangle\langle\, v_i \,|$, via the square roots of the eigenvalues λ_i. Then $\sqrt{f} \circ \sqrt{f} = f$ because the eigenvectors v_i are pairwise orthogonal. The 'square root lemma' says that for each positive operator f on a Hilbert space there is a unique positive $\sqrt{f}$ with $\sqrt{f} \circ \sqrt{f} = f$, see *e.g.* [29, Theorem 1.33]. If f is a projection one has $\sqrt{f} = f$, because a projection's eigenvalues are either 0 or 1.

The following result is essential.

Lemma 15. *For a positive predicate $p \colon X \to X \oplus X$ in* ***FdHilb****, the above map $\sqrt{p} \colon X \to X \oplus X$ is a dagger mono.*

As a result there is a substitution functor $(\sqrt{p})^* \colon \mathit{pPred}(X \oplus X) \to \mathit{pPred}(X)$ as in Proposition 6. It satisfies:

$$p \;=\; \big(\sqrt{p}\big)^*(\Omega_X),$$

so that we can take $\mathrm{char}_p = \sqrt{p}$ in this quantum case.

Proof. If $p = \langle p_1, p_2 \rangle \colon X \to X \oplus X$ is a positive predicate, then p_i is positive, and so is $\sqrt{p_i}$; in particular, $\sqrt{p_i}$ is self-adjoint. Hence we simply compute:

$$\begin{aligned}
(\sqrt{p})^\dagger \circ \sqrt{p} &= \langle \sqrt{p_1}, \sqrt{p_2} \rangle^\dagger \circ \langle \sqrt{p_1}, \sqrt{p_2} \rangle \\
&= [\sqrt{p_1}^\dagger, \sqrt{p_2}^\dagger] \circ \langle \sqrt{p_1}, \sqrt{p_2} \rangle \\
&= [\sqrt{p_1}, \sqrt{p_2}] \circ \langle \sqrt{p_1}, \sqrt{p_2} \rangle \\
&= \nabla \circ (\sqrt{p_1} \oplus \sqrt{p_2}) \circ (\sqrt{p_1} \oplus \sqrt{p_2}) \circ \Delta \\
&= \nabla \circ (p_1 \oplus p_2) \circ \Delta \\
&= \nabla \circ \langle p_1, p_2 \rangle \\
&= \mathrm{id}.
\end{aligned}$$

This map $\sqrt{p}\colon X \rightarrowtail X \oplus X$ in †-*Mono*(**Hilb**) satisfies, using substitution as described in (10):

$$\begin{aligned}
\big(\sqrt{p}\big)^*(\varOmega_X) \;&=\; (\sqrt{p}^\dagger \oplus \sqrt{p}^\dagger) \circ \varOmega_X \circ \sqrt{p} \\
&=\; ([\sqrt{p_1}, \sqrt{p_2}] \oplus [\sqrt{p_1}, \sqrt{p_2}]) \circ (\kappa_1 \oplus \kappa_2) \circ \langle \sqrt{p_1}, \sqrt{p_2} \rangle \\
&=\; (\sqrt{p_1} \oplus \sqrt{p_2}) \circ \langle \sqrt{p_1}, \sqrt{p_2} \rangle \\
&=\; \langle \sqrt{p_1} \circ \sqrt{p_1}, \sqrt{p_2} \circ \sqrt{p_2} \rangle \\
&=\; \langle p_1, p_2 \rangle \\
&=\; p. \qquad\qquad \square
\end{aligned}$$

Also in this quantum case there is no uniqueness, since also the pair $\langle -\sqrt{p_1}, -\sqrt{p_2} \rangle$ can serve as characteristic map. But if we require that char_p consists of positive maps, then the pair $\langle \sqrt{p_1}, \sqrt{p_2} \rangle$ is the only possible choice.

9.1 Towards dynamic logic

Characteristic maps will be used in measurement, in the next section. But they can also be used to define generic test operators $\langle p? \rangle$ and $[p?]$ like in dynamic logic [28].

Definition 8. *Let $p, q\colon X \to X+X$ be two predicates in a category with characteristic maps char and substitution. We define:*

$$\begin{aligned}
\langle p? \rangle(q) \;&=\; \big(\mathit{char}_p\big)^*\big[(\kappa_1 + \kappa_1) \circ q, \kappa_2 \circ \kappa_2\big] \\
[p?](q) \;&=\; \big(\mathit{char}_p\big)^*\big[(\kappa_1 + \kappa_1) \circ q, \kappa_1 \circ \kappa_2\big].
\end{aligned}$$

The cotuples $[\cdots]$ on the right hand side involve different ways of extending q from a predicate on X to a predicate $X + X \to (X + X) + (X + X)$ on $X + X$.

Below we briefly check what these operations are in our three main examples, and leave further investigation of the logical rules involved (like in [5]) to future work. The informal reading of the predicate $\langle p? \rangle(q)$ is as “p andthen q”. This is in general not a commutative operation, see the third example below (in Hilbert spaces). In the other two examples, this “andthen” is the same as the effect monoid multiplication from Example 15. The common reading of $[p?](q)$ is “p then q”, understood as a weak version of $\langle p? \rangle(q)$. We briefly use these operations in Example 19.

Example 18. In the predicate logic of sets, predicates $X \to X + X$ can be identified with subsets $U \subseteq X$, see Example 1. In terms of these subsets the formulations in Definition 8 translate into the familiar conjunction and implications for subsets:

$$\begin{aligned}
\langle U? \rangle(V) \;&=\; U \cap V \\
[U?](V) \;&=\; U \Rightarrow V \;=\; \neg(U \cap \neg V).
\end{aligned}$$

Predicates in the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of the distribution monad are in fact fuzzy predicates, see Example 3. For two such predicates $\varphi, \psi \in [0,1]^X$ on X we have:

$$\begin{aligned}
\langle \varphi ? \rangle(\psi)(x) &= \varphi(x) \cdot \psi(x) \\
[\varphi ?](\psi)(x) &= \varphi(x) \cdot \psi(x) + 1 - \varphi(x) \\
&= 1 - \varphi(x) \cdot (1 - \psi(x)) \;=\; \big(\varphi(x) \cdot \psi(x)^{\perp}\big)^{\perp}.
\end{aligned}$$

This last formula gives the so-called Reichenbach implication [54].

Finally, in Hilbert spaces we can identify predicates on a space X with effects on X, see Example 5. For two such effects $A, B \colon X \to X$ Definition 8 yields:

$$\begin{aligned}
\langle A ? \rangle(B) &= \sqrt{A}\, B \sqrt{A} \\
[A ?](B) &= \big(\sqrt{A}\, B \sqrt{A}\big) + \mathrm{id} - A.
\end{aligned}$$

As far as we know these formulas are new in "dynamic quantum logic". We illustrate their use in Example 19.

10 Measurement

Measurement involves two key aspects, namely probability calculation, and passage to successor states. In the present context the former is done via substitution, see Example 13; the latter will be described below, via the characteristic maps introduced in the previous section.

At this stage we assume we have a category **C** with an indexed category $P \colon \mathbf{C}^{\mathrm{op}} \to \mathbf{EMod}$ for predicates. For convenience we leave the underlying effect monoid of probabilities $\mathrm{Pr}(\mathbf{C})$ often implicit. The main examples involve the categories **Sets**, $\mathcal{K}\ell(\mathcal{D})$, and **FdHilb**.

Given the indexed category P of predicates we define a new *states* functor $S \colon \mathbf{C} \to \mathbf{Conv}$, where **Conv** is the category of convex sets, over the effect monoid $\mathrm{Pr}(\mathbf{C})$, see Subsection 5.3.

$$\begin{aligned}
S(X) &= \mathbf{EMod}\big(P(X), \mathrm{Pr}(\mathbf{C})\big) \\
S\big(X \xrightarrow{f} Y\big) &= - \circ f^{*} \;:\; \mathbf{EMod}\big(P(X), \mathrm{Pr}(\mathbf{C})\big) \longrightarrow \mathbf{EMod}\big(P(Y), \mathrm{Pr}(\mathbf{C})\big).
\end{aligned}$$

A state $s \in S(X)$ is thus a function that transforms predicates on X into probabilities (predicates on the tensor unit). This makes a state a predicate transformer, as used in classical work on program semantics, see [18]. Explicitly, $s(p) \in \mathrm{Pr}(\mathbf{C})$ is the probability that predicate p holds for s.

Using the adjunction from Proposition 2 we may describe the situation as follows.

$$\begin{tikzcd}
& \mathbf{EMod}^{\mathrm{op}} \arrow[rr, bend left, "Hom(-{,}\mathrm{Pr}(\mathbf{C}))"] \arrow[rr, phantom, "\top"] & & \mathbf{Conv} \arrow[ll, bend left, "Hom(-{,}\mathrm{Pr}(\mathbf{C}))"] & \\
\text{predicates} & & \mathbf{C} \arrow[ul, "P"] \arrow[ur, "S"'] & & \text{states}
\end{tikzcd}$$

The left (predicate) part of this diagram is associated with Heisenberg, and the right (state) part with Schrödinger, see [29].

Measurement is specified here by a predicate $p\colon X \to X + X$. Measurement then exists in three forms, namely for pure states, for states as predicate-transformers, and for mixed states. Measurement wrt. p uses the associated characteristic map $\mathit{char}_p\colon X \to X + X$ in the base category of the indexed category.

(1) For pure states, that is, for elements $x\colon I \to X$ in the base category, *pure state* measurement yields the pure state $\mathit{char}_p \circ x\colon I \to X + X$.
(2) For states as predicate-transformers, measurement is the map:

$$S(X) \xrightarrow{S(\mathit{char}_p)} S(X + X)$$

(3) Mixed states $M(X)$ arise for certain objects X as alternative descriptions $M(X) \cong S(X)$ of predicates-to-probabilities. Via this isomorphism one gets mixed state measurement as map $M(X) \to M(X + X)$.

This description gives only a pattern, which will be elaborated for our three main examples in the remainder of this section. In each case the coproduct type $X + X$ for measured states gives a pattern-match outcome, making it possible to proceed differently, depending on whether the predicate used in measurement holds or not: these coproducts "represent the branching arising from the indeterminacy of measurement outcomes" (cited from [2, §§6, point 6a]). Such a coproduct $X + X$ is the appropriate output type, because only by taking both outcomes (left and right) into account one gets a probability distribution, with sums of probabilities adding up to 1[3].

Set-theoretic measurement In the set-theoretic case we have booleans $2 = \{0, 1\}$ as probabilities. The predicate functor is powerset $\mathcal{P}\colon \mathbf{Sets}^{\mathrm{op}} \to \mathbf{EMod}_2 = \mathbf{EA}$. The associated state functor $S\colon \mathbf{Sets} \to \mathbf{Conv}_2 = \mathbf{Sets}$ is:

$$S(X) = \mathbf{EA}\big(\mathcal{P}(X), 2\big) = \mathbf{BA}\big(\mathcal{P}(X), 2\big), \qquad \text{by Lemma 8.}$$

This S is also known as the *ultrafilter* monad (see [46,39]).

For a predicate $p\colon X \to X + X$ in **Sets** the pure state measurement map is p itself:

$$\mathit{Hom}(1, X) \cong X \xrightarrow{\mathit{char}_p = p} X + X \cong \mathit{Hom}(1, X + X)$$

Measurement on predicate transformers is described by:

$$S(X) = \mathbf{BA}\big(\mathcal{P}(X), 2\big) \xrightarrow{S(\mathit{char}_p) = S(p) = (-) \circ p^*} \mathbf{BA}\big(\mathcal{P}(X + X), 2\big) = S(X + X)$$

---

[3] Instead of using such coproduct outcomes the probability or trace is often normalised via a suitable division, like in $\frac{P\rho P}{tr(\rho P)}$ for a density matrix ρ. But this may fail, typically if one has to divide by 0. In the present situation this handled by adding $P^{\perp}$ in the right part of a coproduct.

where p^* is the substitution functor

$$\mathit{Pred}(X+X) = \mathcal{P}(X+X) \xrightarrow{\;p^*\;} \mathcal{P}(X) = \mathit{Pred}(X)$$

If X is a finite set, then by the finitary version of Stone duality (see *e.g.* [4]) one has:

$$S(X) \;=\; \mathbf{BA}\big(\mathcal{P}(X), 2\big) \;\cong\; X.$$

Thus pure states, mixed states and states as predicate-transformers are the all same in the (finite) set-theoretic case. Via this isomorphism the measurement map $S(\mathit{char}_p)$ is simply the predicate p itself, measurement operation:

$$X \xrightarrow{\;p\;} X+X$$

This predicate p, as measurement operation, sends a state $x \in X$ to either the left or the right component of $X+X$, depending on whether p holds for x. We shall see a similar behaviour in the other examples.

Probabilistic measurement We now work in the Kleisli category $\mathcal{K}\ell(\mathcal{D})$ of sets and stochastic relations. The predicates $\mathit{Pred}(X)$ on a set X can be described as fuzzy relations $[0,1]^X$. As a result, the state functor $S\colon \mathbf{Sets} \to \mathbf{Conv}$ is given by:

$$S(X) \;=\; \mathbf{EMod}\big([0,1]^X, [0,1]\big).$$

This is the *expectation* monad from [37].

Assume a predicate $p\colon X \to X+X$ in the category $\mathcal{K}\ell(\mathcal{D})$. A pure state $m\colon 1 \to X$ corresponds to a distribution $m \in \mathcal{D}(X)$. The pure measurement operation is thus given by Kleisli composition:

$$\mathit{Hom}(1,X) \cong \mathcal{D}(X) \xrightarrow{\;\mathit{char}_p \circ (-) = p \circ (-)\;} \mathcal{D}(X+X) \cong \mathit{Hom}(1, X+X)$$

This map is the Kleisli extension $p^{\#}$ of the function $p\colon X \to \mathcal{D}(X+X)$, see Subsection 2.3 and (17) below.

Measurement on predicate-transformers involves the map:

$$\begin{tikzcd}
\mathbf{EMod}\big([0,1]^X, [0,1]\big) \arrow[r, "S(\mathit{char}_p) = S(p) = (-)\circ p^*"] \arrow[d, phantom, "\parallel"] & \mathbf{EMod}\big([0,1]^{X+X}, [0,1]\big) \arrow[d, phantom, "\parallel"] \\
S(X) & S(X+X)
\end{tikzcd}$$

involving the substitution functor:

$$\mathit{Pred}(X+X) = [0,1]^{X+X} \xrightarrow{\;p^*\;} [0,1]^X = \mathit{Pred}(X),$$

given by $p^*(\psi)(x) = p(x)(\kappa_1 x)\cdot\psi(\kappa_1 x) + p(x)(\kappa_2 x)\cdot\psi(\kappa_2 x)$.

As noted in [37], if X is a finite set, there is an isomorphism of convex sets, between the distribution monad and the expectation monad on X:

$$\mathcal{D}(X) \xrightarrow[\cong]{\xi} \mathbf{EMod}\big([0,1]^X, [0,1]\big), \tag{16}$$

given by $\xi(m)(\varphi) = \sum_x m(x)\cdot\varphi(x)$. Thus the mixed states are the distributions $\mathcal{D}(X)$, and coincide with the pure states and with the states as predicate-transformer, also in the probabilistic case. Via the isomorphism ξ we get a mixed-state measurement operation which coincides with pure-state measurement:

$$\mathcal{D}(X) \xrightarrow{p^{\#}} \mathcal{D}(X+X) \quad \text{given by} \quad \begin{cases} p^{\#}(m)(\kappa_1 x) \;=\; m(x)\cdot p(x)(\kappa_1 x) \\ p^{\#}(m)(\kappa_2 x) \;=\; m(x)\cdot p(x)(\kappa_2 x) \end{cases} \tag{17}$$

A distribution $m \in \mathcal{D}(X)$ yields via $p^{\#}$ a new distribution in $\mathcal{D}(X+X)$, where the probability of an element $\kappa_1 x$ in the first component is given by m's probability of x multiplied by the probability that the predicate p holds for x. Similarly, for $\kappa_2 x$, one uses the probability $p(x)(\kappa_2 x) = 1 - p(x)(\kappa_1 x)$ that p does *not* hold for x. Thus, the predicate p splits the state m into two parts, as a state with type $X+X$, and multiplies the probabilities in m with p's probabilities, as weights.

Quantum measurement for Hilbert spaces We now assume a positive predicate $p = \langle p_1, p_2\rangle \colon H \to H \oplus H$ on a finite-dimensional Hilbert space H, where $p_1 \colon H \to H$ is an effect with orthocomplement $p_2 = \mathrm{id} - p_1 = p_1^{\perp}$. There is an associated dagger mono $\mathit{char}_p = \sqrt{p} = \langle \sqrt{p_1}, \sqrt{p_2}\rangle \colon H \rightarrowtail H \oplus H$, as described before Lemma 15.

The base category is the subcategory †-*Mono*(**FdHilb**) of dagger monos. Measurement of pure states happens simply via post-composition on dagger monos:

$$\Big(I \xrightarrow{x} X\Big) \longmapsto \Big(I \xrightarrow{\mathit{char}_p \circ x} X \oplus X\Big) \tag{18}$$

Thus, an element $x \in H$ with $\|x\|^2 = 1$ is sent to the successor state $x' = \mathit{char}_p \circ x = \sqrt{p} \circ x \in H \oplus H$, also with $\|x'\|^2 = 1$. This successor state consists of two parts, $\sqrt{p_1} \circ x$ on the left with appropriate amplitudes incorporating the probability that the measurement wrt. predicate p succeeds, and $\sqrt{p_2} \circ x$ on the right where it does not succeed. This is elaborated in an example below.

On states as predicate-transformers we get the map:

$$\begin{tikzcd}
\mathbf{EMod}\big(\mathcal{E}f(H),[0,1]\big) \arrow[r, "S(\mathit{char}_p)=S(\sqrt{p})=(-)\circ(\sqrt{p})^*"] \arrow[d, equal] & \mathbf{EMod}\big(\mathcal{E}f(H\oplus H),[0,1]\big) \arrow[d, equal] \\
S(H) & S(H\oplus H)
\end{tikzcd} \tag{19}$$

where we use that $\sqrt{p}$ is a dagger mono, and so, by Proposition 6, we have a substitution functor $(\sqrt{p})^* \colon \mathcal{E}f(H \oplus H) \cong \mathit{pPred}(H+H) \to \mathit{pPred}(H) \cong \mathcal{E}f(H)$, given by $A \mapsto (\sqrt{p})^{\dagger} \circ A \circ \sqrt{p}$, following (10).

For mixed state measurement we use that state-transformers can be described as density matrices (positive operators with trace 1), via the isomorphism,

$$\mathcal{DM}(H) \xrightarrow[\cong]{\xi} \mathbf{EMod}\big(\mathcal{E}f(H), [0,1]\big),$$

given by $\xi(\rho)(A) = tr(\rho A)$, see [9,38]. Via this ξ we obtain the following measurement operation for mixed states.

$$\begin{array}{rcl} \mathcal{DM}(H) & \longrightarrow & \mathcal{DM}(H \oplus H) \\ \rho & \longmapsto & \sqrt{p} \circ \rho \circ (\sqrt{p})^\dagger = \langle \sqrt{p_1}, \sqrt{p_2} \rangle \circ \rho \circ [\sqrt{p_1}, \sqrt{p_2}]. \end{array} \tag{20}$$

For convenience, write $\rho' = \sqrt{p} \circ \rho \circ (\sqrt{p})^\dagger \colon H \oplus H \to H \oplus H$. This ρ' is again a density matrix, since:

- ρ' is positive: if $\rho = f^\dagger \circ f$, then $g = f \circ (\sqrt{p})^\dagger$ yields $g^\dagger \circ g = \rho'$;
- ρ' has trace 1, since by the cyclic property of the trace operation, that is, $tr(AB) = tr(BA)$, one has:

$$\begin{array}{rcll} tr(\rho') \;=\; tr(\sqrt{p} \circ \rho \circ (\sqrt{p})^\dagger) & = & tr(\rho \circ (\sqrt{p})^\dagger \circ \sqrt{p}) & \\ & = & tr(\rho) & \text{since } \sqrt{p} \text{ is a dagger mono} \\ & = & 1. & \end{array}$$

In order to check that predicate-transformer measurement (19) and mixed-state measurement (20) coincide we consider the following diagram.

$$\begin{tikzcd} \mathcal{DM}(H) \arrow[r, "\xi", "\cong"'] \arrow[d, "\sqrt{p}\circ(-)\circ(\sqrt{p})^\dagger"'] & \mathbf{EMod}\big(\mathcal{E}f(H), [0,1]\big) \arrow[d, "S(\sqrt{p})=(-)\circ(\sqrt{p})^*"] \\ \mathcal{DM}(H \oplus H) \arrow[r, "\xi", "\cong"'] & \mathbf{EMod}\big(\mathcal{E}f(H \oplus H), [0,1]\big) \end{tikzcd}$$

It commutes, since for $\rho \in \mathcal{DM}(H)$ and $A \in \mathcal{E}f(H \oplus H)$,

$$\begin{array}{rcll} \big(S(\sqrt{p}) \circ \xi\big)(\rho)(A) & = & S(\sqrt{p})(\xi(\rho))(A) & \\ & = & \big(\xi(\rho) \circ (\sqrt{p})^*\big)(A) & \\ & = & \xi(\rho)((\sqrt{p})^\dagger \circ A \circ \sqrt{p}) & \text{by (10)} \\ & = & tr(\rho \circ (\sqrt{p})^\dagger \circ A \circ \sqrt{p}) & \\ & = & tr(\sqrt{p} \circ \rho \circ (\sqrt{p})^\dagger \circ A) & \text{by the cyclic property} \\ & = & \xi(\sqrt{p} \circ \rho \circ (\sqrt{p})^\dagger)(A) & \\ & = & \big(\xi \circ (\sqrt{p} \circ (-) \circ (\sqrt{p})^\dagger)\big)(\rho)(A). & \end{array}$$

Finally, if we apply mixed-state measurement $\mathcal{DM}(H) \to \mathcal{DM}(H \oplus H)$ in (20) to a density matrix $x \circ x^\dagger$ coming from a pure state $x \colon I \to X$, then the outcome is the density matrix coming from the result $x' = \sqrt{p} \circ x$ of pure-state measurement in (18), since:

$$x' \circ x'^{\dagger} \,=\, (\sqrt{p} \circ x) \circ (\sqrt{p} \circ x)^{\dagger} \,=\, \sqrt{p} \circ (x \circ x^{\dagger}) \circ (\sqrt{p})^{\dagger}.$$

In the end we like to offer an explanation for the occurrence of the square root $\sqrt{p}$ of predicates in quantum measurement. Assume the associated effect $p_1 = \pi_1 \circ p$ has spectral decomposition $\sum_i \lambda_i |\, v_i \,\rangle\langle\, v_i \,|$, with eigenvalues $\lambda_i \in [0,1]$ and orthonormal eigenvectors $|\, v_i \,\rangle$. These λ_i are weights, or multiplication factors, for the probability of having a measurement outcome $|\, v_i \,\rangle$, as illustrated in Example 13. The probabilities of the measurement outcomes are obtained, via the Born rule, as squares of the absolute values of the amplitudes. Hence in order to multiply these probabilities with λ_i, one has to multiply the underlying amplitudes with $\sqrt{\lambda_i}$. This happens in measurement wrt. to a predicate p, namely via $\sqrt{p}$. Thus, eigenvalues are not irrelevant in measurement (as argued *e.g.* in [55]): they do have a precise role when we use measurements specified by predicates (*i.e.* effects). The categorical approach used here gives eigenvalues their proper role, namely as probability scalars.

Remark 5. As illustrated above, predicates $p = \langle p_1, p_2 \rangle \colon X \to X \oplus X$ can be used for "binary" measurement. It is quite natural to do n-ary measurement, say via a map $p = \langle p_1, \ldots, p_n \rangle \colon X \to X \oplus \cdots \oplus X$, where each p_i is an effect, and $\sum_i p_i = \mathrm{id}$.

In case we have an n-ary *projective* measurement, these p_i's are projections, which are pairwise orthogonal — so that $p_i \circ p_j = 0$ for $i \neq j$. Such projective measurement can be reduced to predicates as used here. This will be illustrated for $n = 3$. So assume we have orthogonal projections $p_1, p_2, p_3 \colon X \to X$ with $p_1 + p_2 + p_3 = \mathrm{id}$. Then we can form the following two predicates on X.

$$q_1 \,=\, \langle p_1, \mathrm{id} - p_1 \rangle \qquad \text{and} \qquad q_2 \,=\, \langle p_2, \mathrm{id} - p_2 \rangle.$$

By consecutively doing q_1 and then q_2 on the second $\oplus$-component we get the triple $\langle p_1, p_2, p_3 \rangle \colon X \to X \oplus X \oplus X$, since:

$$\begin{aligned}
(\mathrm{id} \oplus q_2) \circ q_1 &= \langle p_1,\ \langle p_2, \mathrm{id} - p_2 \rangle \circ (\mathrm{id} - p_1) \rangle \\
&= \langle p_1,\ \langle p_2 \circ (\mathrm{id} - p_1),\ (\mathrm{id} - p_2) \circ (\mathrm{id} - p_1) \rangle \rangle \\
&= \langle p_1,\ \langle p_2 - (p_2 \circ p_1),\ \mathrm{id} - p_2 - ((\mathrm{id} - p_2) \circ p_1) \rangle \rangle \\
&= \langle p_1,\ \langle p_2 - 0,\ \mathrm{id} - p_2 - (p_1 - (p_2 \circ p_1)) \rangle \rangle \\
&= \langle p_1,\ \langle p_2,\ \mathrm{id} - p_2 - p_1 + 0 \rangle \rangle \\
&= \langle p_1,\ \langle p_2,\ p_3 \rangle \rangle.
\end{aligned}$$

The more general form of measurement described in [51, §§2.2.3] can be reduced to such projective measurement, see *loc. cit.*

We conclude with a probability calculation in the present setting for the famous polarisation experiment, described for instance in [55].

Example 19. The idea in the polarisation experiment is to send photons polarised as $|\uparrow\rangle = |\, 0 \,\rangle = \left(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right)$ through filters.

(1) If these photons hit one filter with polarisation $|\,\rightarrow\,\rangle = |\,1\,\rangle = \left(\begin{smallmatrix}0\\1\end{smallmatrix}\right)$, then nothing passes.

(2) Next assume there are two consecutive filters: first the $|\,0\,\rangle$-photons have to go through a filter with polarisation $|\,\nearrow\,\rangle = \frac{1}{\sqrt{2}}\left(\begin{smallmatrix}1\\1\end{smallmatrix}\right)$, and then through a filter with polarisation $|\,\rightarrow\,\rangle = |\,1\,\rangle = \left(\begin{smallmatrix}0\\1\end{smallmatrix}\right)$. Surprisingly, now one quarter of the original photons get through. We refer for an explanation to *e.g.* [55], and focus on a calculation of this probability $\frac{1}{4}$ in the current quantitative logic.

For the experiment (1) we use the predicate $p_{|\,1\,\rangle} \colon \mathbb{C}^2 \rightarrow \mathbb{C}^2 \oplus \mathbb{C}^2$ given by:

$$p_{|\,1\,\rangle} \;=\; \langle\, |1\rangle\langle\, 1\,|,\ \mathrm{id} - |1\rangle\langle\, 1\,|\,\rangle \;=\; \langle\, |1\rangle\langle\, 1\,|,\ |0\rangle\langle\, 0\,|\,\rangle \;=\; \langle\, \left(\begin{smallmatrix}0&0\\0&1\end{smallmatrix}\right),\ \left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)\rangle$$

The probability of a photon $|\,0\,\rangle$ getting through filter $|\,1\,\rangle$ is obtained via the quantum substitution formula (10) (or Born rule), like in Example 13:

$$\begin{aligned}
\pi_1 \circ |\,0\,\rangle^*(p_{|\,1\,\rangle}) &= \pi_1 \circ (|\,0\,\rangle^\dagger \oplus |\,0\,\rangle^\dagger) \circ p_{|\,1\,\rangle} \circ |\,0\,\rangle \\
&= |\,0\,\rangle^\dagger \circ \pi_1 \circ p_{|\,1\,\rangle} \circ |\,0\,\rangle \\
&= \langle 0\,|\,1\rangle\langle 1\,|\,0\rangle \\
&= 0.
\end{aligned}$$

For experiment (2), with an intermediate $|\,\nearrow\,\rangle$-filter, we use the associated predicate $p_{|\,\nearrow\,\rangle} \colon \mathbb{C}^2 \rightarrow \mathbb{C}^2 \oplus \mathbb{C}^2$ given by:

$$\begin{aligned}
p_{|\,\nearrow\,\rangle} &= \langle\, |\,\nearrow\,\rangle\langle\,\nearrow\,|,\ \mathrm{id} - |\,\nearrow\,\rangle\langle\,\nearrow\,|\,\rangle \;=\; \langle\, |\,\nearrow\,\rangle\langle\,\nearrow\,|,\ |\,\nwarrow\,\rangle\langle\,\nwarrow\,|\,\rangle \\
&= \langle\, \tfrac{1}{2}\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right),\ \tfrac{1}{2}\left(\begin{smallmatrix}1&-1\\-1&1\end{smallmatrix}\right)\rangle,
\end{aligned}$$

where $|\,\nwarrow\,\rangle = \frac{1}{\sqrt{2}}\left(\begin{smallmatrix}1\\-1\end{smallmatrix}\right)$. This predicate satisfies $\sqrt{p_{|\,\nearrow\,\rangle}} = p_{|\,\nearrow\,\rangle}$ since it is a projection. The probability that a $|\,0\,\rangle$-polarised photon gets through a $|\,\nearrow\,\rangle$-filter is given, as above, by:

$$\pi_1 \circ |\,0\,\rangle^*(p_{|\nearrow\rangle}) \;=\; |\,0\,\rangle^\dagger|\,\nearrow\,\rangle\langle\,\nearrow\,||0\rangle \;=\; \langle 0\,|\,\nearrow\rangle\langle 0\,|\,\nearrow\rangle \;=\; \tfrac{1}{\sqrt{2}} \cdot \tfrac{1}{\sqrt{2}} \;=\; \tfrac{1}{2}.$$

But we are interested in the consecutive use of two filters, $p_{|\,1\,\rangle}$ after $p_{|\,\nearrow\,\rangle}$. We shall use the "$p_{|\,\nearrow\,\rangle}$ andthen $p_{|\,1\,\rangle}$" dynamic operator $\langle p_{|\,\nearrow\,\rangle}?\rangle(p_{|\,1\,\rangle})$ from Subsection 9.1 for this purpose. As effect it can be described like in Example 18 as:

$$\begin{aligned}
\big\langle p_{|\,\nearrow\,\rangle}?\big\rangle(p_{|\,1\,\rangle}) &= \sqrt{\pi_1 \circ p_{|\,\nearrow\,\rangle}}(\pi_1 \circ p_{|\,1\,\rangle})\sqrt{\pi_1 \circ p_{|\,\nearrow\,\rangle}} \\
&= |\,\nearrow\,\rangle\langle\,\nearrow\,|\,|1\rangle\langle\, 1\,|\,|\,\nearrow\,\rangle\langle\,\nearrow\,| \\
&= \tfrac{1}{2}\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right)\left(\begin{smallmatrix}0&0\\0&1\end{smallmatrix}\right)\tfrac{1}{2}\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right) \\
&= \tfrac{1}{4}\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right).
\end{aligned}$$

The probability of $\frac{1}{4}$ in the above experiment (2) is now calculated via substitution (Born) as follows.

$$\begin{aligned}
|\,0\,\rangle^*\big\langle p_{|\,\nearrow\,\rangle}?\big\rangle(p_{|\,1\,\rangle}) &= |\,0\,\rangle^\dagger\big\langle p_{|\,\nearrow\,\rangle}?\big\rangle(p_{|\,1\,\rangle})|0\rangle \\
&= \left(\begin{smallmatrix}1&0\end{smallmatrix}\right)\tfrac{1}{4}\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right)\left(\begin{smallmatrix}1\\0\end{smallmatrix}\right) \;=\; \tfrac{1}{4}\left(\begin{smallmatrix}1&0\end{smallmatrix}\right)\left(\begin{smallmatrix}1\\1\end{smallmatrix}\right) \;=\; \tfrac{1}{4}.
\end{aligned}$$

As an afterthought we add that the weaker "$p_{|\nearrow\rangle}$ then $p_{|1\rangle}$" test operator $[p_{|\nearrow\rangle}?](p_{|1\rangle})$ adds the option that the photon does *not* get through the first filter $|\nearrow\rangle$ — which happens with chance $\frac{1}{2}$ — and thus yields as outcome:

$$|0\rangle^{*}\big[p_{|\nearrow\rangle}?\big](p_{|1\rangle}) \,=\, \tfrac{3}{4}.$$

Details of this calculation, via the corresponding formula for effects in Example 18, are left to the reader.

11 Conclusions

This paper has elaborated the first steps of a new categorical logic based on predicates as maps of the form $X \to X + X$. It has shown that this alternative representation makes sense, especially in a quantitative setting, for classical, probabilistic, and quantum logic; it involves operations in the style of dynamic logic and measurement wrt. a predicate in each of these logics, via appropriate characteristic maps.

The current work offers several avenues for further research. As already mentioned in the introduction, tensors do not really play a role yet in the current framework, and their logic is still missing. Such extension is needed for a comparison between the tensor-based approach to measurement "without sums" in [15], and the current logical/quantitative approach in which these sums are fundamental (in order to define predicates in the first place). Also, it would be interesting what the "strange" aspects of quantum theory, like non-locality and contextuality, look like in the current logical setting. On the categorical side, a precise axiomatisation of the structures used in this paper may lead to further clarity (through abstraction), and possibly to a notion of a "quantitative topos". On the syntactical side a precise description of predicates, their effect module structure, substitution, and test operations $[p?]$ and $\langle p?\rangle$, together with a proof calculus, could be useful.

Acknowledgements Thanks go to Jorik Mandemaker, Chris Heunen, Robert Furber, and Ichiro Hasuo for helpful discussions and feedback.

References

1. S. Abramsky and B. Coecke. A categorical semantics of quantum protocols. In *Logic in Computer Science*, pages 415–425. IEEE, Computer Science Press, 2004.
2. S. Abramsky and B. Coecke. A categorical semantics of quantum protocols. In K. Engesser, Dov M. Gabbai, and D. Lehmann, editors, *Handbook of Quantum Logic and Quantum Structures: Quantum Logic*, pages 261–323. North Holland, Elsevier, Computer Science Press, 2009.
3. M.A. Arbib and E.G. Manes. *Algebraic Approaches to Program Semantics*. Texts and Monogr. in Comp. Sci. Springer, Berlin, 1986.
4. S. Awodey. *Category Theory*. Oxford Logic Guides. Oxford Univ. Press, 2006.
5. A. Baltag and S. Smets. LQP: the dynamic logic of quantum information. *Math. Struct. in Comp. Sci.*, 16:491–525, 2006.

6. G. Birkhoff and J. von Neumann. The logic of quantum mechanics. *Ann. Math.*, 37:823–843, 1936.
7. F. Borceux. *Handbook of Categorical Algebra*, volume 50, 51 and 52 of *Encyclopedia of Mathematics*. Cambridge Univ. Press, 1994.
8. M. Born. Quantenmechanik der stoßvorgänge. *Zeitschrift f. Physik*, 38:803–827, 1926.
9. P. Busch. Quantum states and generalized observables: a simple proof of Gleason's Theorem. *Phys. Review Letters*, 91(12):120403:1–4, 2003.
10. J. Butterfield and C.J. Isham. A topos perspective on the Kochen-Specker theorem: I. quantum states as generalized valuations. *Int. Journ. Theor. Physics*, 37(11):2669–2733, 1998.
11. A. Carboni, S. Lack, and R.F.C. Walters. Introduction to extensive and distributive categories. *Journ. of Pure & Appl. Algebra*, 84(2):145–158, 1993.
12. M. Caspers, C. Heunen, K. Landsman, and B. Spitters. Intuitionistic quantum logic of an n-level system. *Found. of Physics*, 39:731–759, 2009.
13. F. Chovanec and F. Kôpka. D-posets. In K. Engesser, Dov M. Gabbai, and D. Lehmann, editors, *Handbook of Quantum Logic and Quantum Structures: Quantum Structures*, pages 367–428. North Holland, Elsevier, Computer Science Press, 2007.
14. J.R.B. Cockett. Introduction to distributive categories. *Math. Struct. in Comp. Sci.*, 3:277–307, 1993.
15. B. Coecke and D. Pavlović. Quantum measurements without sums. In G. Chen, L. Kauffman, and S. Lamonaco, editors, *Mathematics of Quantum Computing and Technology*. Taylor and Francis, 2006. See also: arxiv:/quant-ph/0608035.
16. E. D'Hondt and P. Panangaden. Quantum weakest preconditions. *Math. Struct. in Comp. Sci.*, 16(3):429–451, 2006.
17. M. Dickson. Aspects of probability in quantum physics. In C. Beisbart and S. Hartmann, editors, *Probabilities in Physics*, pages 171–199. Oxford Univ. Press, New York, 2011.
18. E.W. Dijkstra and C. Scholten. *Predicate Calculus and Program Semantics*. Springer, Berlin, 1990.
19. A. Döring and C.J. Isham. A topos foundation for theories of physics: I - IV. *Journ. of Math. Physics*, 49:053515–053518, 2008.
20. A. Dvurečenskij and S. Pulmannová. *New Trends in Quantum Structures*. Kluwer Acad. Publ., Dordrecht, 2000.
21. D. J. Foulis and M.K. Bennett. Effect algebras and unsharp quantum logics. *Found. Physics*, 24(10):1331–1352, 1994.
22. D. J. Foulis, R.J. Greechie, and M.K. Bennett. The transition to unigroups. *Int. Journ. Theor. Physics*, 37(1):45–63, 1998.
23. R. Giuntini and H. Greuling. Toward a formal language for unsharp properties. *Found. Physics*, 19:769–780, 1994.
24. A. Gleason. Measures on the closed subspaces of a Hilbert space. *Journ. Math. Mech.*, 6:885–893, 1957.
25. S. Gudder. Examples, problems, and results in effect algebras. *Int. Journ. Theor. Physics*, 35(11):2365–2376, 1996.
26. E. Haghverdi. Unique decomposition categories, geometry of interaction and combinatory logic. *Math. Struct. in Comp. Sci.*, 10:205–231, 2000.
27. P. Halmos. *Introduction to Hilbert Space and the Theory of spectral Multiplicity*. Chelsea Publ. Comp., New York, 2^{nd} edition, 1957.
28. D Harel, D. Kozen, and J. Tiuryn. *Dynamic Logic*. MIT Press, Cambridge, MA, 2000.
29. T. Heinosaari and M. Ziman. *The Mathematical Language of Quantum Theory. From Uncertainty to Entanglement*. Cambridge Univ. Press, 2012.
30. C. Heunen and B. Jacobs. Quantum logic in dagger kernel categories. *Order*, 27(2):177–212, 2010.

31. C. Heunen, N.P. Landsman, and B. Spitters. A topos for algebraic quantum theory. *Comm. in Math. Physics*, 291:63–110, 2009.
32. B. Jacobs. *Categorical Logic and Type Theory*. North Holland, Amsterdam, 1999.
33. B. Jacobs. Convexity, duality, and effects. In C.S. Calude and V. Sassone, editors, *IFIP Theoretical Computer Science 2010*, number 82(1) in IFIP Adv. in Inf. and Comm. Techn., pages 1–19. Springer, Boston, 2010.
34. B. Jacobs. Coalgebraic walks, in quantum and Turing computation. In M. Hofmann, editor, *Foundations of Software Science and Computation Structures*, number 6604 in Lect. Notes Comp. Sci., pages 12–26. Springer, Berlin, 2011.
35. B. Jacobs. Probabilities, distribution monads, and convex categories. *Theor. Comp. Sci.*, 412(28):3323–3336, 2011.
36. B. Jacobs and J. Mandemaker. Coreflections in algebraic quantum logic. *Found. of Physics*, 10 May 2012. `http://dx.doi.org/doi:10.1007/s10701-012-9654-8`.
37. B. Jacobs and J. Mandemaker. The expectation monad in quantum foundations. In B. Jacobs, P. Selinger, and B. Spitters, editors, *Quantum Physics and Logic (QPL) 2011*, 2012. EPTCS, to appear; see arxiv.org/abs/1112.3805.
38. B. Jacobs and J. Mandemaker. Relating operator spaces via adjunctions. In J. Chubb Reimann, V. Harizanov, and A. Eskandarian, editors, *Logic and Algebraic Structures in Quantum Computing and Information*, Lect. Notes in Logic. Cambridge Univ. Press, 2012. See arxiv.org/abs/1201.1272.
39. P.T. Johnstone. *Stone Spaces*. Number 3 in Cambridge Studies in Advanced Mathematics. Cambridge Univ. Press, 1982.
40. G. Kalmbach. *Orthomodular Lattices*. Academic Press, London, 1983.
41. G.M. Kelly and M. L. Laplaza. Coherence for compact closed categories. *Journ. of Pure & Appl. Algebra*, 19:193–213, 1980.
42. S. Kochen and E. Specker. The problem of hidden variables in quantum mechanics. *J. of Math. & Mech.*, 17:59–87, 1967.
43. K. Kraus. *States, Effects, and Operations*. Springer Verlag, Berlin, 1983.
44. N.P. Landsman. Born rule and its interpretation. In D. Greenberger, K. Hentschel, and F. Weinert, editors, *Compendium of Quantum Physics*, pages 64–70. Springer, 2009.
45. G. Ludwig. *Foundations of Quantum Mechanics I*. Springer Verlag, New York, 1983.
46. E.G. Manes. A triple-theoretic construction of compact algebras. In B. Eckman, editor, *Seminar on Triples and Categorical Homolgy Theory*, number 80 in Lect. Notes Math., pages 91–118. Springer, Berlin, 1969.
47. P. Martin-Löf. *Intuitionistic Type Theory*. Bibliopolis, Napoli, 1984.
48. S. Mac Lane. *Categories for the Working Mathematician*. Springer, Berlin, 1971.
49. S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic. A First Introduction to Topos Theory*. Springer, New York, 1992.
50. J. von Neumann. *Mathematische Grundlagen der Quantenmechanik*. Springer, Berlin, 1932.
51. M.A. Nielsen and I.L. Chuang. *Quantum Computation and Quantum Information*. Cambridge Univ. Press, 2000.
52. B. Nordström, K. Peterson, and J.M. Smith. *Programming in Martin-Löf's Type Theory: an introduction*. Number 7 in Logic Guides. Oxford Science Publ., 1990.
53. S. Pulmannová and S. Gudder. Representation theorem for convex effect algebras. *Commentationes Mathematicae Universitatis Carolinae*, 39(4):645–659, 1998. Available from `http://dml.cz/dmlcz/119041`.
54. H. Reichenbach. *The theory of probability*. Univ. California Press, 1949.
55. E. Rieffel and W. Polak. *Quantum Computing. A Gentle Introduction*. MIT Press, Cambridge, MA, 2011.
56. M.H. Stone. Postulates for the barycentric calculus. *Ann. Math.*, 29:25–30, 1949.

57. T. Swirszcz. Monadic functors and convexity. *Bull. de l'Acad. Polonaise des Sciences. Sér. des sciences math., astr. et phys.*, 22:39–42, 1974.
58. S. Venegas-Andraca. *Quantum Walks for Computer Scientists*. Morgan & Claypool, 2008.